

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07K 14/195, 16/12, C12N 1/21, 15/31, G01N 33/569, A61K 39/02, 39/40		A1	(11) International Publication Number: WO 97/36923 (43) International Publication Date: 9 October 1997 (09.10.97)
(21) International Application Number: PCT/AU97/00212 (22) International Filing Date: 1 April 1997 (01.04.97)		(81) Designated States: AU, CA, JP, KR, NZ, SG, US, European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).	
(30) Priority Data: PN 9112 29 March 1996 (29.03.96) AU		Published With international search report.	
(71) Applicants (for all designated States except US): THE UNIVERSITY OF MELBOURNE [AU/AU]; Grattan Street, Parkville, VIC 3052 (AU). VICTORIAN DAIRY INDUSTRY AUTHORITY [AU/AU]; 651-653 Victoria Street, Abbotsford, VIC 3067 (AU).			
(72) Inventors; and (75) Inventors/Applicants (for US only): REYNOLDS, Eric, Charles [AU/AU]; 104 Hill Street, North Balwyn, VIC 3104 (AU). SLAKESKI, Nada [AU/AU]; 4 Findon Crescent, Kew, VIC 3101 (AU). HENDTLASS, Anne [GB/AU]; 26 Grosvenor Parade, Balwyn, VIC 3103 (AU).			
(74) Agent: F.B. RICE & CO.; 28A Montague Street, Balmain, NSW 2041 (AU).			

(54) Title: PORPHYROMONAS GINGIVALIS ANTIGENS FOR THE DIAGNOSIS AND TREATMENT OF PERIODONTITIS

(57) Abstract

The present invention provides a composition for use in raising an immune response directed against *Porphyromonas gingivalis*. The composition includes a suitable adjuvant and/or acceptable carrier and one substantially purified *P. gingivalis* immunogen. The immunogen is selected from the group consisting of Antigen 1, Antigen 2, Antigen 3, Antigen 4 and epitope containing fragments thereof, in which: Antigen 1 is an antigen of *P. gingivalis* and has an internal amino acid sequence: DLENKGEATLLVTFGSSYKAPRETYAKIEKTFAAAYPDQR; Antigen 2 is an antigen of *P. gingivalis* and has an internal amino acid sequence: DNPDENPLEGDIQTHTTEKYVLAED; Antigen 3 is an antigen of *P. gingivalis* and has an internal amino acid sequence: DVLLLDVTPLSLGIETMGGVMTYLIDANTTPKLLK; Antigen 4 is an antigen of *P. gingivalis* and has an internal amino acid sequence: VYNASISAVGNTSAIDPVVQIIHHN.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TK	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NI	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroun	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

Porphyromonas gingivalis Antigens for the Diagnosis and Treatment
of Periodontitis

FIELD OF THE INVENTION

This invention relates to oral compositions and immunogenic compositions
5 for use in the suppression of the pathogenic effects of the bacterium
Porphyromonas gingivalis associated with periodontal disease and
cardiovascular disease. It also relates to diagnostic tests for the presence of
Porphyromonas gingivalis in subgingival plaque samples and specific
10 antibodies against *P. gingivalis* antigens. The compositions comprise
proteins, peptides or oligopeptides or peptide chimeras of specific antigens
of *Porphyromonas gingivalis*. Also disclosed are methods for preparing the
antigens, peptide components and peptide chimeras using recombinant DNA
and/or biochemical techniques. Related thereto, disclosed are the DNA
15 sequences encoding the specific antigens, and recombinant vectors useful in
directing the expression of antigen constructs containing major epitopes.
Also disclosed are host cells transformed with such recombinant vectors.
The proteins, peptides, oligopeptides and peptide chimeras are useful as
immunogens in formulations for use in raising an immune response and can
20 be used to generate protein-specific and peptide-specific antisera useful for
passive immunization and as reagents for diagnostic assays. The nucleotide
sequences disclosed provide for the synthesis of corresponding
oligonucleotides which can be used as reagents in diagnostic assays directed
to the detection of *P. gingivalis* genetic material and incorporated into
expression vectors for use as genetic vaccine formulations.

25 BACKGROUND OF THE INVENTION

Periodontal diseases are bacterial-associated inflammatory diseases of the
supporting tissues of the teeth and range from the relatively mild form of
gingivitis, the non-specific, reversible inflammation of gingival tissue to the
more aggressive forms of periodontitis which are characterised by the
30 destruction of the tooth's supporting structures. Periodontitis is associated
with a subgingival infection of a consortium of specific Gram-negative
bacteria that leads to the destruction of the periodontium and is a major
public health problem. One bacterium that has attracted considerable
interest is *Porphyromonas gingivalis* as the recovery of this microorganism
35 from adult periodontitis lesions can be up to 50% of the subgingival
anaerobically cultivable flora, whereas *P. gingivalis* is rarely recovered, and

then in low numbers, from healthy sites. A proportional increase in the level of *P. gingivalis* in subgingival plaque has been associated with an increased severity of periodontitis and eradication of the microorganism from the cultivable subgingival microbial population is accompanied by resolution of the disease. The progression of periodontitis lesions in non-human primates has been demonstrated with the subgingival implantation of *P. gingivalis*. These findings in both animals and humans suggest a major role for *P. gingivalis* in the development of adult periodontitis. The presence of *P. gingivalis* in atheromatous plaques has also been associated with the development of cardiovascular disease.

P. gingivalis is a black-pigmented, anaerobic, proteolytic Gram-negative rod that obtains energy from the metabolism of specific amino acids. The microorganism has an absolute growth requirement for iron, preferentially in the form of heme or its Fe(III) oxidation product hemin and when grown under conditions of excess hemin is highly virulent in experimental animals. A number of virulence factors have been implicated in the pathogenicity of *P. gingivalis* including the capsule, adhesins, cytotoxins and extracellular hydrolytic enzymes. In order to develop an efficacious and safe vaccine to prevent *P. gingivalis* colonisation it is necessary to identify protein antigens that are involved in virulence that have utility as immunogens to generate neutralising antibodies.

SUMMARY OF THE INVENTION

The present inventors purified and characterised four major *P. gingivalis* antigens using serum from a healthy subject that harboured *P. gingivalis* in subgingival plaque as shown by DNA probe analysis. The antigens (Ag1, Ag2, Ag3 and Ag4) are listed below

<u>Ag</u>	<u>M_r</u>	<u>Putative Function</u>	<u>Internal Amino acid sequence</u>
Ag1	32kDa	Haeme receptor	DLENKGEATLLVTFGSSYKAPRETYAKIEK TFAAAYPDQR
Ag2	46kDa	Fimbrial protein	DNPDENPLEGDITQTHTEKYVLAED
Ag3	70kDa	DnaK homologue	DVLLLDVTPLSLGIETMGGVMTYLIDANTT IPKLK
Ag4	10kDa	S-layer protein	VYNASISAVGNTSAIDPVVQIIHHN (N-terminal sequence)

Accordingly in a first aspect the present invention consists in a composition for use in raising an immune response directed against *Porphyromonas gingivalis*, the composition including a suitable adjuvant and/or acceptable carrier and one substantially purified *P. gingivalis* immunogen, the immunogen being selected from the group consisting of Antigen 1, Antigen 2, Antigen 3, Antigen 4 and epitope containing fragments thereof.

Optionally, the composition may further include at least one additional purified *P. gingivalis* immunogen, the immunogen being selected from the group consisting of Antigen 1, Antigen 2, Antigen 3, Antigen 4 and epitope containing fragments thereof.

In a second aspect, the present invention consists in a substantially purified *P. gingivalis* antigen or epitope containing fragment thereof, wherein antigen has an internal amino acid sequence:

DLENKGEATLLVTFGSSYKAPRETYAKIEKTFAAAYPDQR It is preferred that the antigen has an amino acid sequence as shown in Fig 1

In a third aspect, the present invention consists in a substantially purified *P. gingivalis* antigen or epitope containing fragment thereof, wherein antigen has an internal amino acid sequence:

DNPDENPLEGDITQ'THTEKYVLAED.

- 5 In a fourth aspect, the present invention consists in a substantially purified *P. gingivalis* antigen or epitope containing fragment thereof, wherein antigen has an internal amino acid sequence:

DVLLLDVTPLSLGIETMGGVMTYLIDANTTIPKLLK. It is preferred that the antigen includes an amino acid sequence encoded by the open reading frame of the clone deposited with AGAL under accession No. NM 97/04974 which hybridises with degenerate probes corresponding to the amino acid sequence DVLLLDVTPLSLGIETMGGVMTYLIDANTTIPKLLK.

- 10

In a fourth aspect, the present invention consists in a substantially purified *P. gingivalis* antigen or epitope containing fragment thereof, wherein antigen has an internal amino acid sequence:

- 15 VYNASISAVGNTSAIDPVVQIIHHN.

In other aspects, the present invention consists in nucleotide sequences encoding Ag1, Ag2, Ag3 and Ag4 and probes which hybridise to these sequences.

- 20 The nucleotide sequence encoding Ag1 and deduced amino acid sequence of the haeme receptor protein Ag1 is shown in Fig 1. The disclosure of the nucleotide sequence includes within its scope degeneracy equivalents and subsequences coding for amino acid sequences corresponding to antigenic determinants of *P. gingivalis* W50.

- 25 A clone containing nucleotide sequence from Ag3, DnaK clone #6, has been deposited under the terms of the Budapest Treaty with Australian Government Analytical Laboratories, 1 Suakin Street, Pyrmble, NSW Australia on 25 March 1997 and has been accorded accession No. NM 97/04974. Accordingly further nucleotide sequence for this antigen can be obtained by accessing this deposit. Access to this deposit is available under the terms and conditions of the Budapest Treaty. Where applicable access to this deposit is to be limited to independent experts (EPC Rule 28(4) A.U. Reg. 3.25(3)).

- 30 In another aspect the present invention consists in antibodies raised against the antigens of the present invention
- 35

Antibodies against the antigens can be used in oral compositions such as toothpaste and mouthwash to neutralise the antigens and thus prevent disease. Antigen-specific antibodies can also be used for the early detection of *P. gingivalis* in subgingival plaque samples by a diagnostic assay. A vaccine based on these antigens and suitable adjuvant delivered by nasal spray, orally or by injection to produce a specific immune response against these antigens thereby reducing colonisation and virulence of *P. gingivalis* and thereby preventing or reducing disease. The antigen proteins and antigen peptides (herein termed "peptides") and antigen oligopeptides (herein termed "oligopeptides") and antigen chimeric peptides containing epitopes of one antigen fused with the epitopes of another (herein termed "chimeric peptides") thereof, of the present invention may be used as immunogens in prophylactic and/or therapeutic vaccine formulations; or as an antigen in diagnostic immunoassays directed to detection of *P. gingivalis* infection by measuring an increase in serum titer of *P. gingivalis* - specific antibody. Also antigen protein, peptides, oligopeptides and chimeric peptides of the present invention may be used to generate antigen-specific antibody which may be useful for passive immunotherapy and as reagents for diagnostic assays directed to detecting the presence of *P. gingivalis* in clinical specimens such as subgingival plaque samples. Peptides, oligopeptides or chimeric peptides can be obtained by chemical synthesis, purification from *P. gingivalis* cultures, or produced from recombinant vector expression systems using the nucleic acid sequences disclosed herein. Accordingly, in other aspects the invention provides oral compositions including toothpastes and mouthwashes which include antibodies raised against any one or a combination of antigens Ag1, Ag2, Ag3 and Ag4. In another aspect the invention provides a method of early detection of *P. gingivalis* comprising a diagnostic assay involving the use of antibodies raised against any one or a combination of antigens Ag1, Ag2, Ag3 and Ag4. In another aspect, the invention provides a method for the detection of *P. gingivalis* infection comprising the measure of an increase in serum titer to any one of the *P. gingivalis* antigens as herein described. Other aspects of the present invention are directed to the construction of novel DNA sequences involving *antigen* constructs and vectors including plasmid DNA, and viral DNA such as human viruses, animal viruses, insect viruses, or bacteriophages which can be used to direct the expression of

antigen protein, peptides, oligopeptides or chimeric peptides in appropriate host cells from which the expressed protein or peptides may be purified. Another aspect of the present invention provides methods for molecular cloning of the genes encoding the antigens Ag1, Ag2, Ag3 and Ag4, and gene fragments encoding antigen peptides or oligopeptides or chimeric peptides. The nucleic acid sequences of the present invention can be used in molecular diagnostic assays for *P. gingivalis* genetic material through nucleic acid hybridization, and including the synthesis of antigen sequence-specific oligonucleotides for use as primers and/or probes in amplifying, and detecting amplified, nucleic acids. Additionally, antigen protein, peptides, oligopeptides, chimeric peptides and antigenic constructs containing epitopes can be used as immunogens in prophylactic and/or therapeutic vaccine formulations against pathogenic strains of *P. gingivalis*, whether the immunogen is chemically synthesized, purified from *P. gingivalis*, or purified from a recombinant expression vector system. Alternatively, the genes encoding the antigens, or one or more gene fragments encoding peptides or oligopeptides or chimeric peptides, may be incorporated into a bacterial or viral vaccine comprising recombinant bacteria or virus which is engineered to produce one or more immunogenic epitopes of each antigen by itself, or in combination with immunogenic epitopes of other antigens or from other pathogenic microorganisms. In addition, the genes encoding the antigens or one or more gene fragments encoding peptides or oligopeptides or chimeric peptides, operatively linked to one or more regulatory elements, can be introduced directly into humans to express protein, peptide, oligopeptides or chimeric peptides relating to the antigens to elicit a protective immune response. A vaccine can also be based upon a recombinant component of a mutated antigen incorporated into an appropriate vector and expressed in a suitable transformed host (e.g. *E. coli*, *Bacillus subtilis*, *Saccharomyces cerevisiae*, COS cells, CHO cells and HeLa cells) containing the vector. The vaccine can be based on an intra-oral recombinant bacterial vaccine, where the recombinant bacterium expressing antigen is a commensal inhabitant of the oral cavity. Unlike whole *P. gingivalis* cells or other previously prepared antigens, the four antigens described herein are safe and effective antigens for the preparation of a vaccine for the prevention of *P. gingivalis*-associated periodontal disease.

In yet another aspect the present consists in a composition for use in raising an immune response directed against *Porphyromonas gingivalis*, the composition including a suitable adjuvant and/or acceptable carrier and a DNA molecule including a sequence encoding one *P. gingivalis* immunogen, the immunogen being selected from the group consisting of Antigen 1, Antigen 2, Antigen 3, Antigen 4 and epitope containing fragments thereof. *P. gingivalis* has an absolute growth requirement for Fe which it prefers in the form of haeme. As such, Ag1 is of particular interest as neutralisation of this haeme receptor by specific antibodies would prevent haeme uptake and therefore growth and virulence. *P. gingivalis* grown haeme-limited is less virulent in animal models.

Fimbriae are thin, filamentous structures that either completely cover the cell or are polar. There are at least two fimbrial types recognized: those involved in the transfer of genetic material by the formation of conjugation bridges, and those involved in adherence to soft and hard tissues. The fimbriae involved in conjugation are referred to as sex pili. These pili have specific receptors for attachment to a genetically compatible recipient bacterium. The second fimbrial type, the type specific or common pili are involved in eubacterial coaggregation and adherence to eukaryotic cells and often play an important role in prevention of phagocytosis and the invasion of host tissue. In the *Enterobacteriaceae*, the fimbriae consist of repeating subunit proteins of approximately 17 to 21 kDa. Minor proteins are also part of the fimbriae structure. The specific fimbrial binding proteins (adhesins) are often 28 to 31 kDa and located at the tip or periodically along the length of the fimbriae.

Yoshimura *et al.* were amongst the first to demonstrate the presence of fimbriae on *P. gingivalis* and they purified a 43 kDa fimbriilin subunit which has no amino acid sequence homology with fimbriilins from other Gram-negative bacteria. Genco and coworkers have shown that the 43 kDa fimbrial protein and synthetic peptides corresponding to the C-terminal end of the fimbriilin reduced adherence of *P. gingivalis* 381 to saliva-coated hydroxyapatite. In a subsequent study they showed that a recombinant fimbriilin binds specifically to statherin and proline-rich proteins of saliva. Immunisation of rats with the 43 kDa protein protected against periodontal tissue destruction induced by infection with *P. gingivalis* 381. Further, an isogenic mutant of *P. gingivalis* 381 with the *fimA* gene, that encodes the 43

kDa fimbrilin, insertionally inactivated was significantly less able to produce periodontal tissue destruction in the rat model when compared with the wild-type strain. In this study it should be noted however, that the *fimA* mutant still did produce greater periodontal tissue destruction than occurred in the sham infected animals. The *fimA* mutant was unimpaired in its ability to agglutinate red blood cells, coaggregate with other oral bacteria although binding to saliva-coated hydroxyapatite was reduced. In an independent study Hamada *et al.* also produced a *fimA* mutant of *P. gingivalis* ATCC 33277 by homologous recombination of an insertionally inactivated gene and noted that although the mutant failed to express long (0.5 to 1.0 μ m) fimbriae, thin, short fimbrial structures could still be observed by electron microscopy suggesting the presence of a second fimbrial type. The identification of the 43 kDa fimbrilin and the virulence and immunisation studies related to the 43 kDa protein have been conducted with the *P. gingivalis* strains 381 and ATCC 33277. These strains are classified as non-invasive and are considered to be less virulent than invasive strains based on the infective process in animal models. Non-invasive strains produce a localised abscess at the challenged site, whereas invasive strains at the same inoculum spread to distant sites and produce multiple abscesses. Further, Sundqvist *et al.* showed that the non-invasive strains (381 and ATCC 33277) were phagocytosed and killed by polymorphonuclear leukocytes to a high extent whereas the invasive strains W50 and W83 were poorly phagocytosed and killed. It is interesting to note that Naito *et al.* suggest that there is a difference in the fimbriae of invasive and non-invasive *P. gingivalis* strains. These workers found that the fimbriae of non-invasive strains bound to collagen-coated hydroxyapatite (HA) in high numbers whereas the fimbriae prepared from invasive strains bound to collagen-coated HA weakly. The *P. gingivalis* invasive strains W50, W83 and AJW5 are highly virulent in animal models but do not express the 43 kDa fimbrilin as shown by immunocytochemistry and Western blot analysis. However, on fine negative staining, W50 and W83 are fimbriated although less densely than other strains. It appears that W50 and W83 possess inactive *fimA* genes accounting for the lack of the 43 kDa fimbrilin however these strains are still virulent and invasive despite lacking the 43 kDa fimbrilin.

Ag2 is the second fimbrial type or a major adhesin of *P. gingivalis*. As part of a study to purify and characterise cell surface protein antigens of *P. gingivalis* W50 we purified a 30 kDa fragment of the 46 kDa fimbrial protein (Ag2) that was seroreactive with serum from a healthy subject that
 5 harboured *P. gingivalis* subgingivally as shown by DNA probe analysis. The internal amino acid sequence of the 30 kDa fragment showed considerable homology (48% identity) to a fimbrial protein of *Dichelobacter* (formerly *Bacteroides*) *nodosus*.

	<i>P. gingivalis</i> 30 kDa fragment	DNPDENPLEGDITQTHTEKYVLAED...
10	<i>D. nodosus</i> fimbrial protein	KGPDANPASGVVGKNDTGKYVLAEL...

The *D. nodosus* fimbrial protein is classified as a type-IV or mPhe pilin which is a common fimbrial type of a group of Gram-negative bacteria including *Bacteroides* spp., *Neisseria gonorrhoeae*, *Neisseria meningitidis*,
 15 *Acinetobacter calcoaceticus*, *Eikenella corrodens*, *Moraxella bovis*, *Moraxella nonliquefaciens* and several species of *Pseudomonas* including *P. aeruginosa* (Elleman, 1988). The *P. gingivalis* 30 kDa fragment exhibits the highest homology with the conserved amino acyl residues of the central domain of the *D. nodosus* A-set or Class I fimbriae (including the serotypes A, B, C, E,
 20 F. and G). A characteristic of the type-IV fimbriae is that they adhere to eukaryotic cells and agglutinate red blood cells. It is interesting to note that the conserved hexapeptide motif -KYVLAE- which is also present in the *P. gingivalis* 46 kDa fimbrial protein has been localised to, or near to, the receptor binding site of gonococcal pili since antisera to these residues
 25 prevents bacterial attachment to eukaryotic cells even by heterologous pili and precipitates larger peptide fragments which bind to eukaryotic cells. It has been suggested that the conserved residues of the central domain when juxtaposed form a cleft which specifically interacts with the carbohydrate moieties of surface glycoproteins of eukaryotic cells.
 30 *D. nodosus* is the aetiological agent of the contagious disease of sheep, interdigital dermatitis or footrot, and the type-IV fimbriae are the major serological and immunoprotective virulence factors. Footrot vaccines have evolved from simple bacterins to highly specific recombinant DNA fimbrial vaccines. The initial whole cell vaccines were unsuccessful due to the short
 35 duration of immunity and incorporation of limited serotypes. A number of antigens were examined and the major protective immunogen was the type-

IV fimbrial subunit protein. Monovalent vaccines based on recombinant fimbriae are omnipotent inducing long lasting immunity. The homology between the *P. gingivalis* 46 kDa fimbrial protein and the *D. nodosus* immunoprotective fimbriae indicate that the *P. gingivalis* protein (Ag2) would have application in diagnostic and immunoprophylactic products for *P. gingivalis*-related periodontitis.

The heat shock or stress response of cells is a homeostatic mechanism that enables cells to survive environmental stresses such as temperature elevation that can result in denaturation of cellular proteins. The DnaK family of proteins bind to denatured and incorrectly folded proteins and facilitate refolding to the original conformation and function. The DnaK or Heat Shock Protein (HSP) 70 is a highly conserved molecular chaperonin common to bacterial and eukaryotic cells comprising 1 - 5% of the constitutive cellular protein, with 15% of DnaK in *E. coli* being associated with vesicles. During stress DnaK can be overexpressed to constitute up to 30% of total cellular protein making this protein an ideal candidate for a sensitive immunodiagnostic test. Specific diagnostic tests for leprosy and tuberculosis have been developed based on the respective DnaK protein homologues. All species homologues of DnaK are highly conserved in the N-terminal half of the protein with the species-specific regions of the molecule in the C-terminal half. The *P. gingivalis* DnaK homologue (Ag3) therefore would have application in diagnostic and immunoprophylactic products for *P. gingivalis*-related periodontitis.

The four antigens identified (Ag1, Ag2, Ag3 and Ag4) are of particular interest for diagnostics and neutralisation by passive immunity through oral compositions containing neutralising antibodies and by vaccine development. In particular for the development of an intra-oral recombinant bacterial vaccine, where the recombinant bacterium expressing the antigens is a genetically engineered commensal inhabitant of the oral cavity. The superiority of these four antigens to prior disclosed *P. gingivalis* antigens, is that these are major virulence-associated factors and contain conserved epitopes on invasive strains making them ideal for the development of diagnostic and immunoprophylactic products.

DETAILED DESCRIPTION OF THE INVENTION

The present invention will now be described in detail with reference to the following particularly preferred embodiments which are not limiting to the

invention but representative of the methods of performing certain aspects of the invention

The four antigens Ag1, Ag2, Ag3 and Ag4 can be purified from *P. gingivalis* cells by chloroform extraction followed by anion exchange, gel filtration and reversed-phase chromatography. The purified antigens are then used to generate polyclonal or monoclonal antibodies using standard techniques. The animals used for antibody generation can be mice, rabbits, goats, chickens, sheep, horses, cows etc. When a high antibody titre against the antigens is detected by immunoassay the animals are bled or eggs or milk are collected and the serum prepared and/or antibody purified using standard techniques or monoclonal antibodies produced by fusing spleen cells with myeloma cells using standard techniques. The antibody (immunoglobulin fraction) may be separated from the culture or ascites fluid, serum, milk or egg by salting out, gel filtration, ion exchange and/or affinity chromatography, and the like, with salting out being preferred. In the salting out method the antiserum or the milk is saturated with ammonium sulphate to produce a precipitate, followed by dialyzing the precipitate against physiological saline to obtain the purified immunoglobulin fraction with the specific antibody. The preferred antibody is obtained from the equine antiserum and the bovine antiserum and milk. In this invention the antibody contained in the antiserum and milk obtained by immunising the animal with the antigens is blended into the oral composition. In this case the antiserum and milk as well as the antibody separated and purified from the antiserum and milk may be used. Each of these materials may be used alone or in combination of two or more. Antibodies can be used in oral compositions such as toothpaste and mouthwash to neutralise *P. gingivalis* and thus prevent disease. The antibodies can also be used for the early detection of *P. gingivalis* in subgingival plaque samples by a chairside Enzyme Linked Immunosorbent Assay (ELISA).

For oral compositions it is preferred that the amount of the above antibodies administered is 0.0001 -50 g/kg/day and that the content of the above antibodies is 0.0002 - 10% by weight preferably 0.002 -5% by weight of the composition. The oral composition of this invention which contains the above-mentioned serum or milk antibody may be prepared and used in various forms applicable to the mouth such as dentifrice including toothpastes, toothpowders and liquid dentifrices, mouthwashes, troches.

chewing gums, dental pastes, gingival massage creams, gargle tablets, dairy products and other foodstuffs. The oral composition according to this invention may further include additional well known ingredients depending on the type and form of a particular oral composition.

- 5 In certain highly preferred forms of the invention the oral composition may be substantially liquid in character, such as a mouthwash or rinse. In such a preparation the vehicle is typically a water-alcohol mixture desirably including a humectant as described below. Generally, the weight ratio of water to alcohol is in the range of from about 1:1 to about 20:1. The total
10 amount of water-alcohol mixture in this type of preparation is typically in the range of from about 70 to about 99.9% by weight of the preparation. The alcohol is typically ethanol or isopropanol. Ethanol is preferred. The pH of such liquid and other preparations of the invention is generally in the range of from about 4.5 to about 9 and typically from about 5.5 to 8. The
15 pH is preferably in the range of from about 6 to about 8.0, preferably 7.4. The pH can be controlled with acid (e.g. citric acid or benzoic acid) or base (e.g. sodium hydroxide) or buffered (as with sodium citrate, benzoate, carbonate, or bicarbonate, disodium hydrogen phosphate, sodium dihydrogen phosphate, etc).
- 20 Other desirable forms of this invention, the oral composition may be substantially solid or pasty in character, such as toothpowder, a dental tablet or a dentifrice, that is a toothpaste (dental cream) or gel dentifrice. The vehicle of such solid or pasty oral preparations generally contains dentally acceptable polishing material. Examples of polishing materials are water-
25 insoluble sodium metaphosphate, potassium metaphosphate, tricalcium phosphate, dihydrated calcium phosphate, anhydrous dicalcium phosphate, calcium pyrophosphate, magnesium orthophosphate, trimagnesium phosphate, calcium carbonate, hydrated alumina, calcined alumina, aluminum silicate, zirconium silicate, silica, bentonite, and mixtures
30 thereof. Other suitable polishing material include the particulate thermosetting resins such as melamine-, phenolic, and urea-formaldehydes, and cross-linked polyepoxides and polyesters. Preferred polishing materials include crystalline silica having particle sized of up to about 5 microns, a mean particle size of up to about 1.1 microns, and a surface area of up to
35 about 50,000 cm²/gm, silica gel or colloidal silica, and complex amorphous alkali metal aluminosilicate.

When visually clear gels are employed, a polishing agent of colloidal silica, such as those sold under the trademark SYLOID as Syloid 72 and Syloid 74 or under the trademark SANTOCEL as Santocel 100, alkali metal aluminosilicate complexes are particularly useful since they have refractive indices
5 close to the refractive indices of gelling agent-liquid (including water and/or humectant) systems commonly used in dentifrices.

Many of the so-called "water insoluble" polishing materials are anionic in character and also include small amounts of soluble material. Thus, insoluble sodium metaphosphate may be formed in any suitable manner as
10 illustrated by Thorpe's Dictionary of Applied Chemistry, Volume 9, 4th Edition, pp. 510-511. The forms of insoluble sodium metaphosphate known as Madrell's salt and Kurrol's salt are further examples of suitable materials. These metaphosphate salts exhibit only a minute solubility in water, and therefore are commonly referred to as insoluble metaphosphates (IMP).

15 There is present therein a minor amount of soluble phosphate material as impurities, usually a few percent such as up to 4% by weight. The amount of soluble phosphate material, which is believed to include a soluble sodium trimetaphosphate in the case of insoluble metaphosphate, may be reduced or eliminated by washing with water if desired. The insoluble alkali metal
20 metaphosphate is typically employed in powder form of a particle size such that no more than 1% of the material is larger than 37 microns.

The polishing material is generally present in the solid or pasty compositions in weight concentrations of about 10% to about 99%. Preferably, it is present in amounts from about 10% to about 75% in
25 toothpaste, and from about 70% to about 99% in toothpowder. In toothpastes, when the polishing material is silicious in nature, it is generally present in amount of about 10-30% by weight. Other polishing materials are typically present in amount of about 30-75% by weight.

In a toothpaste, the liquid vehicle may comprise water and humectant
30 typically in an amount ranging from about 10% to about 80% by weight of the preparation. Glycerine, propylene glycol, sorbitol and polypropylene glycol exemplify suitable humectants/carriers. Also advantageous are liquid mixtures of water, glycerine and sorbitol. In clear gels where the refractive index is an important consideration, about 2.5 - 30% w/w of water, 0 to about
35 70% w/w of glycerine and about 20-80% w/w of sorbitol are preferably employed

- Toothpaste, creams and gels typically contain a natural or synthetic thickener or gelling agent in proportions of about 0.1 to about 10, preferably about 0.5 to about 5% w/w. A suitable thickener is synthetic hectorite, a synthetic colloidal magnesium alkali metal silicate complex clay available for example as Laponite (e.g. CP, SP 2002, D) marketed by Laporte Industries Limited. Laponite D is, approximately by weight 58.00% SiO_2 , 25.40% MgO , 3.05% Na_2O , 0.98% Li_2O , and some water and trace metals. Its true specific gravity is 2.53 and it has an apparent bulk density of 1.0 g/ml at 8% moisture.
- Other suitable thickeners include Irish moss, iota carrageenan, gum tragacanth, starch, polyvinylpyrrolidone, hydroxyethylpropylcellulose, hydroxybutyl methyl cellulose, hydroxypropyl methyl cellulose, hydroxyethyl cellulose (e.g. available as Natrosol), sodium carboxymethyl cellulose, and colloidal silica such as finely ground Syloid (e.g. 244).
- Solubilizing agents may also be included such as humectant polyols such as propylene glycol, dipropylene glycol and hexylene glycol, cellosolves such as methyl cellosolve and ethyl cellosolve, vegetable oils and waxes containing at least about 12 carbons in a straight chain such as olive oil, castor oil and petrolatum and esters such as amyl acetate, ethyl acetate and benzyl benzoate.
- It will be understood that, as is conventional, the oral preparations are to be sold or otherwise distributed in suitable labelled packages. Thus, a jar of mouthrinse will have a label describing it, in substance, as a mouthrinse or mouthwash and having directions for its use; and a toothpaste, cream or gel will usually be in a collapsible tube, typically aluminium, lined lead or plastic, or other squeeze, pump or pressurized dispenser for metering out the contents, having a label describing it, in substance, as a toothpaste, gel or dental cream.
- Organic surface-active agents are used in the compositions of the present invention to achieve increased prophylactic action, assist in achieving thorough and complete dispersion of the active agent throughout the oral cavity, and render the instant compositions more cosmetically acceptable. The organic surface-active material is preferably anionic, nonionic or ampholytic in nature which does not denature the antibody of the invention, and it is preferred to employ as the surface-active agent a detergent material which imparts to the composition detergent and foaming properties while

not denaturing the antibody. Suitable examples of anionic surfactants are water-soluble salts of higher fatty acid monoglyceride monosulfates, such as the sodium salt of the monosulfated monoglyceride of hydrogenated coconut oil fatty acids, higher alkyl sulfates such as sodium lauryl sulfate, alkyl aryl sulfonates such as sodium dodecyl benzene sulfonate, higher alkylsulfoacetates, higher fatty acid esters of 1,2-dihydroxy propane sulfonate, and the substantially saturated higher aliphatic acyl amides of lower aliphatic amino carboxylic acid compounds, such as those having 12 to 16 carbons in the fatty acid, alkyl or acyl radicals, and the like. Examples of the last mentioned amides are N-lauroyl sarcosine, and the sodium, potassium, and ethanolamine salts of N-lauroyl, N-myristoyl, or N-palmitoyl sarcosine which should be substantially free from soap or similar higher fatty acid material. The use of these sarconite compounds in the oral compositions of the present invention is particularly advantageous since these materials exhibit a prolonged marked effect in the inhibition of acid formation in the oral cavity due to carbohydrates breakdown in addition to exerting some reduction in the solubility of tooth enamel in acid solutions. Examples of water-soluble nonionic surfactants suitable for use with antibodies are condensation products of ethylene oxide with various reactive hydrogen-containing compounds reactive therewith having long hydrophobic chains (e.g. aliphatic chains of about 12 to 20 carbon atoms), which condensation products ("ethoxamers") contain hydrophilic polyoxyethylene moieties, such as condensation products of poly (ethylene oxide) with fatty acids, fatty alcohols, fatty amides, polyhydric alcohols (e.g. sorbitan monostearate) and polypropyleneoxide (e.g. Pluronic materials).

Surface active agent is typically present in amount of about 0.1-5% by weight. It is noteworthy, that the surface active agent may assist in the dissolving of the antibody of the invention and thereby diminish the amount of solubilizing humectant needed.

Various other materials may be incorporated in the oral preparations of this invention such as whitening agents, preservatives, silicones, chlorophyll compounds and/or ammoniated material such as urea, diammonium phosphate, and mixtures thereof. These adjuvants, where present, are incorporated in the preparations in amounts which do not substantially adversely affect the properties and characteristics desired.

Any suitable flavoring or sweetening material may also be employed. Examples of suitable flavoring constituents are flavoring oils, e.g. oil of spearmint, peppermint, wintergreen, sassafras, clove, sage, eucalyptus, marjorani, cinnamon, lemon, and orange, and methyl salicylate. Suitable sweetening agents include sucrose, lactose, maltose, sorbitol, xylitol, sodium cyclamate, perillartine, AMP (aspartyl phenyl alanine, methyl ester), saccharine, and the like. Suitably, flavor and sweetening agents may each or together comprise from about 0.1% to 5% more of the preparation.

In the preferred practice of this invention an oral composition according to this invention such as mouthwash or dentifrice containing the composition of the present invention is preferably applied regularly to the gums and teeth, such as every day or every second or third day or preferably from 1 to 3 times daily, at a pH of about 4.5 to about 9, generally about 5.5 to about 8, preferably about 6 to 8, for at least 2 weeks up to 8 weeks or more up to a lifetime.

The compositions of this invention can be incorporated in lozenges, or in chewing gum or other products, e.g. by stirring into a warm gum base or coating the outer surface of a gum base, illustrative of which may be mentioned jelutong, rubber latex, vinylite resins, etc., desirably with conventional plasticizers or softeners, sugar or other sweeteners or such as glucose, sorbitol and the like.

Another important form of the invention is a composition for use in raising an immune response directed against *P. gingivalis* based on the four antigens and suitable adjuvant and/or carrier. This may be delivered via a number of routes, for example by nasal spray, orally or by injection to produce a specific immune response against the antigen thereby reducing colonisation of *P. gingivalis* and reducing virulence thereby preventing disease. As will be readily understood the composition may be based upon a recombinant antigen incorporated into an appropriate vector and expressed in a suitable transformed host (e.g. *E. coli*, *Bacillus subtilis*, *Saccharomyces cerevisiae*, COS cells, CHO cells and HeLa cells) containing the vector. Unlike whole *P. gingivalis* cells or other previously prepared antigens, the antigens described herein or peptides, oligopeptides or chimeric peptides are safe and effective antigens for the preparation of a vaccine for the prevention of *P. gingivalis*-associated periodontal disease. The antigenic protein, peptides, oligopeptides and chimeric peptides of the present invention, can be

- produced using recombinant DNA methods as illustrated herein, or can be synthesized chemically from the amino acid sequence disclosed in the present invention. Additionally, peptides can be produced from enzymatic or chemical cleavage of the purified antigens. Antigenic protein, peptides, and oligopeptides with immunogenic epitopes combined, can be used as immunogens in various vaccine formulations in the prevention of periodontal diseases. Additionally, according to the present invention, antigenic protein and related peptides or chimeras produced may be used to generate *P. gingivalis* antisera useful for passive immunization against periodontal disease and infections caused by *P. gingivalis*.
- As opposed to use of the antigens themselves in eliciting an immune response this may be achieved by administration of a DNA molecule including a sequence encoding at one of the antigens or epitope containing fragment(s).
- The present invention further provides the nucleotide sequence of the genes encoding the antigens, as well as the amino acid sequence deduced from the isolated genes. According to one particularly preferred embodiment of the present invention, using recombinant DNA techniques the genes encoding the antigens or gene fragments encoding one or more peptides or chimeras having immunogenic epitopes, is incorporated into an expression vector, and the recombinant vector is introduced into an appropriate host cell thereby directing the expression of these sequences in that particular host cell. The expression system, comprising the recombinant vector introduced into the host cell, can be used (a) to produce antigenic protein, related peptides, oligopeptides or chimeras which can be purified for use as an immunogen in vaccine formulations; (b) to produce antigenic protein, related peptides, oligopeptides and chimeras to be used as an antigen for diagnostic immunoassays or for generating *P. gingivalis*-specific antisera of therapeutic and/or diagnostic value; (c) or if the recombinant expression vector is a live virus such as vaccinia virus, the vector itself may be used as a live or inactivated vaccine preparation to be introduced into the host's cells for expression of antigen or immunogenic peptides or oligopeptides or chimeric peptides; (d) for introduction into live attenuated bacterial cells or genetically engineered commensal intra-oral bacteria which are used to express antigenic protein, related peptides or oligopeptides or chimeras to vaccinate individuals; (e) or for introduction directly into an individual to

immunize against the encoded and expressed antigenic protein, related peptides, or oligopeptides or chimeras. In particular the recombinant bacterial vaccine can be based on a commensal inhabitant of the human oral cavity or animal if the vaccine is to prevent periodontal disease in animals.

- 5 The recombinant bacterial vaccine expressing antigen can be used to colonise the oral cavity, supragingival or subgingival plaque. The intra-oral bacterium can be isolated from the patient with periodontitis and genetically engineered to express the antigen, peptides or chimeras. The production of the *P. gingivalis* antigen within the oral cavity will not be toxic to the oral
10 mucosal tissues. However, the expressed antigen will stimulate the mucosal-associated lymphoid tissues (MALT) to produce specific antibody to neutralise and reduce the virulence of *P. gingivalis*.

- The following examples are further illustrative of the nature of the present invention, but it is understood that the invention is not limited thereto. All
15 amounts and proportions referred to herein and in the appended claims are by weight unless otherwise indicated.

EXAMPLE 1.

20 Preparation of Antigen 1

Extraction and Purification of Ag1

- P. gingivalis* cells were harvested at O.D._{650nm} 0.18 by centrifugation at 5,000 x g for 20 min at 4°C. The cell pellet was incubated with 10ml
25 chloroform per L of original cell culture with gentle rocking at room temperature for 15 min together with 50ml Buffer A (50 mM NaCl, 10 mM Tris, 10 mM EDTA, pH 8.0) per litre of original culture. The sample was centrifuged at 6,000 x g for 20 min to separate the phases. The aqueous phase was removed and centrifuged a further 30 min at 10,000 x g to
30 eliminate contaminating cell debris. The supernatant was passed through a 0.22µm filter before being applied to a Q sepharose anion exchange chromatography column (XK16/10) at 4°C using an FPLCTM System, (Pharmacia). Proteins were eluted from the column using a linear NaCl gradient from 50 mM NaCl to 500 mM NaCl in 10 mM Tris, 10 mM EDTA,
35 pH 8.0 at 2ml/min over 180 min. Elution was monitored at 280 nm. Six mL fractions were collected and these were analysed by SDS-PAGE and ELISA

using a healthy subject's serum to assess their antigenicity. The most antigenic fraction was concentrated to 200 μ l using a 10,000 NMWCO centricon concentrator before being applied to a Superose 12 (Pharmacia) size-exclusion column.

- 5 The size exclusion column was equilibrated and separation was performed in 150 mM NaCl, 20 mM Tris, 10 mM EDTA, pH 8.0 at 0.3 mL/min. Protein standards were run to enable size comparison with resulting sample peaks. Elution was monitored at 280 nm. Five hundred μ l fractions were collected and these were analysed by SDS-PAGE and ELISA. A peak eluting at 60 kDa
10 was found to be antigenic with serum from a subject harbouring *P. gingivalis* but showing no signs of periodontitis. This fraction was further purified by reversed-phase chromatography.

- Reverse-phase chromatography (RP-HPLC) was performed using a C₁₈RP300 column (Applied Biosystems) and a linear gradient of 0%-80% acetonitrile in
15 0.1% TFA over 40 min. A single peak eluted which contained a single band on SDS-PAGE analysis that had an Mr of 28 kDa. This peak was characterised by N-terminal sequence analysis using a Hewlett Packard 1005A automated protein sequencer.

- From sequence analysis this antigen is likely to be a Haeme receptor of *P. gingivalis* and has an N-terminal amino acid sequence:
20 DLENKGEATLLVTFGSSYKAPRETYAKIEKTFAAAYPDQR:

SDS-PAGE Analysis

- SDS-PAGE analysis was performed using a discontinuous system with a 12%
25 separating gel and a 4% stacking gel. Sample buffer included 4% SDS, and 70 mM β -Mercaptoethanol. Protein standards (Pharmacia) were included on each gel for size comparison. The samples were run at constant voltage of 160 volts in running buffer until the dye-front was near the bottom of the stacking gel. The proteins were visualised with coomassie brilliant blue R-
30 250ml, and destained in 40% MeOH, 7% ILAC in water.

Western Blot analysis

- Western blots were performed to identify proteins which were recognised by serum antibodies from a periodontally diseased (D50) and healthy (H10)
35 subject who harboured *P. gingivalis* subgingivally as shown by DNA analysis. Proteins to be blotted were separated according to the SDS-PAGE procedure.

above, except that pre-stained standards (BioRad) were included to enable the determination of completion of transfer. Following SDS-PAGE, the gel apparatus was dismantled and the gel, PVDF membrane (ProBlott, Applied Biosystems), blotting paper and fibre pads were equilibrated in transfer
5 buffer (10% CAPS, 10% MeOH) for 5 mins. The proteins within the SDS-PAGE gel were transferred onto the PVDF membrane at a constant voltage of 60 volts for 90 min. After transfer the PVDF membrane was blocked with 5% skim milk powder at room temperature for 1 hr. The membrane was then
10 incubated with the primary antibody (human serum H10 1/10 or D50 1/50 diluted in TN buffer (25 mM Tris, 0.5 M NaCl, pH 7.5)) at 4°C overnight. The membrane was washed 3 times for 10 min each in TN buffer then incubated with the second antibody (Horseradish peroxidase conjugated goat anti-human IgG, 1/1000 diluted in TN buffer) at room temperature for
15 2 hours. Once again the membrane was washed 3 times in TN buffer for 10 min each. Binding of the goat anti-human IgG to the antigenic proteins was visualised with 6mg 4-chloro-1-naphthol, 2mL MeOH, 10mL TN buffer and 6µl H₂O₂. The enzyme reaction was stopped with water.

The Western blot analysis showed that the 28 kDa antigen was recognised
20 by the healthy subject (H10) but not by the patient suffering from periodontitis (D50).

ELISA

ELISAs were performed using the serum antibodies from a periodontally
25 diseased (D50) and healthy (H10) subject who harboured *P. gingivalis* subgingivally as shown by DNA analysis. Microtitre plates were coated with the 28 kDa protein or 2% w/v skim milk powder as a negative control, diluted in 50 mM Tris, 200 mM NaCl, pH 7.4 (TBS). The plates were blocked by adding TBS containing 0.05% w/v Tween-20 (TBST) and
30 2% w/v skim milk powder. The wells were washed three times with TBST then incubated with human serum diluted 1/500 in TBST containing 2% w/v polyvinyl-pyrrolidone-40 and 1% w/v NGS (Antibody Diluting Buffer, ADB) for 2 hours at room temperature. The wells were washed three
35 times with TBST then incubated with goat anti-human IgG-horseradish peroxidase diluted 1/3000 in ADB for 2 hours at room temperature. Binding of the anti-human IgG-HRP to the *P. gingivalis* cell-surface proteins was

visualised with 10 mg/mL 3,3',4,4'-tetramethylbenzidine dissolved in dimethyl sulphoxide diluted 1:100 with 0.1 M sodium acetate/citric acid buffer pH 6.0 and 0.004% w/v H₂O₂. The enzyme reaction was stopped with 2 M H₂SO₄ and the colour intensity of the reaction product was
5 spectrophotometrically quantitated by measurement of absorbance at 450 nm.

The ELISA confirmed that the 28 kDa antigen was recognised by the healthy subject (H10) harbouring *P. gingivalis* but not by the patient
10 suffering from periodontitis (D50).

EXAMPLE 2.

15 Cloning and sequence analysis of the Ag1 gene

A lambda GEM 12 library of *Bam*HI digested *P. gingivalis* genomic DNA was screened using degenerate oligonucleotide probes derived from the N-terminal sequence of the 28 kDa antigen (Ag1). A probe-positive clone was
20 identified and contained an insert of 4.6 kbp of *P. gingivalis* genomic DNA. This insert was purified by phenol extraction and was subcloned into pUC18. The DNA sequence of the gene revealed that the open reading frame encoded a protein of predicted mass 32.709. The sequence was determined using the Sanger dideoxy method and the nucleotide sequence and deduced
25 amino acid sequence of Ag1 is presented in Fig 1. The deduced amino acid sequence gives a protein of 32.709 in molecular mass.

EXAMPLE 3.

Construction of a DNA vaccine using the Ag1 gene

30

The plasmid containing the 4.6 kbp of *P. gingivalis* DNA was used as the template for amplification of the gene encoding the 32 kDa antigen (Ag1) by PCR. PCR was performed using the specific oligonucleotide primers with sequences as follows:

35

N-terminal

Oligo A 5'CAA GCA ACA ACA AGG ATT TGC 3'

C-terminal

Oligo B 5'TTG CAT ATC CGC CCG TCC 3'

The PCR mixture contained: Template DNA 100 ng, MgCl₂ 1 mM, Oligo A:800 ng, Oligo B:800 ng, 250 mM dNTP, 3U Ultra polymerase (Perkin Eliner) The thermal cycle involved : Denaturation: 94°C, 30s, Annealing: 52°C, 30s, and Extension: 70°C, 60s.

Southern blot analysis using specific oligonucleotide probes confirmed the correct amplicon. This band was purified from a 1.0% agarose gel by phenol extraction. The amplified fragment was blunt-end ligated with *Sma*I into pUC18 and sequence analysis confirmed the presence of the gene (*Ag1*) encoding the full length sequence of antigen 1. The gene was then subcloned into pcDNA3 (Invitrogen). The clone contained the full length *Ag1* gene (949 base pairs). Approximately 2µg of purified pcDNA3 (Invitrogen) plasmid and DNA from the pUC18 clone 2.2 was digested with *Hind*III (Promega) and *Eco*RI (Promega) restriction enzymes in Multicore buffer (Promega) at 37°C for 1 hour. Digested plasmid DNA was run on 1% Nusieve low melting agarose and the *P. gingivalis* insert DNA cut from the gel and purified using a Bresaclean kit (Bresatech) according to the manufacturer's instructions. The purified insert DNA was resuspended in 20µl of TE buffer. The purified insert DNA from the pUC clones was ligated into the cut pcDNA vector by conventional techniques using T4 ligase (Promega) and ligated at room temperature for 2 hours. Ligated pcDNA3 was transformed into 'Top10F' *E. coli* cells (Invitrogen) which were plated onto agar plates containing ampicillin. Resultant clones were expanded into 3ml cultures, plasmids were purified and screened for insertion of the appropriate sized fragment by restriction enzyme digestion and electrophoresis on 1% agarose gels. Clones showing the correct insert size were selected and further restriction enzyme digests prepared to confirm the insert was correct. This consisted of digesting the clones with the restriction enzymes *Hind*III alone, *Hind*III and *Eco*RI together and *Nde*I alone. One clone was selected 5.1 containing the full length *Ag1* gene. The clone was confirmed by partial DNA sequence analysis using the Sp6 and T7 universal primers following conventional techniques. The identity of the pcDNA3-*Ag1* construct in clone 5.1 was

confirmed by DNA sequence analysis of the 5' and 3' termini, which were as follows:

N-terminus

5' GCT CCC GGC ATC CGC TTA CAG ACA AGC TGT GAC GTC TCC GGG
5 AGC TGC ATG TGT CAG AGG TTT TCA CCG TCA CCG AAA CGC GCG
AGG CTG ATC GTC AGT CAG TCA CGA TGC GGC CGT TCG AGT CGA
CTC TAG AGG ATC CCC CAA GCA ACA ACA AGG ATT TGG AGA ACA
AAG GGG AGG CTA CTC TTT TGG TAA CGT TTG GTA GCT CCT ATA
AAG CTC CA 3'

10 C-terminus

5' GAT GTG TCA AAG ATA TCT GTT CGA CCT GTT ACC GTT AAA GAG
TCG AGA ACC TCT ATA TCT TCT CTA GGC CTT ATA GAC CGA GTT
CGT ATA CTT TCG GTG GAG ACG AGC GAT TCC TGC CCG CCT ATA
CGT TCC CAT GGC TCG AGC TTA AGG ACC CCT AGG TGC GCT TGG
15 TCT AGG CTA AAA CCT CCT ACC AGC GGT GGT GGT TTG CAC GAA 3'

Bolded bases = insert

Non-bolded bases = vector

Large scale (1L) preparations of plasmid DNA was prepared from the clone
20 5.1 and also from the pcDNA3 without any insert. Qiagen Mega Preps were
used to purify the plasmid DNA from overnight cultures of Top 10F⁺ *E. coli*
cells grown in Terrific broth with 150µg/ml of ampicillin. Purified plasmid
DNA was checked by restriction-enzyme analysis to confirm the insert size
and the purity of the DNA. The concentration of DNA/RNA was calculated
25 spectrophotometrically by determining the absorption at 280nm/260nm.

Purified plasmid DNA from the pcDNA3-Ag1 clone with insert (5.1) and
pcDNA3 (no insert) was diluted into 0.9% NaCl in sterile water to 0.5mg/ml
of DNA for injection into animals.

30

EXAMPLE 4.

The ability of the Ag1 DNA vaccine construct to protect against
P. gingivalis challenge in the murine lesion model.

35 10 mice received the pcDNA3-Ag1 construct containing the Ag1 gene
described above (test group), an other 10 mice received pcDNA3 vector

- (control group). 25µg of DNA was injected into each tibialis muscle (a total of 50µg DNA/mouse) of the 20 female, 6 weeks old, Balb/c mice. This was followed a month later with boosts of 25µg of DNA into each tibialis muscle. The mice were subcutaneously challenged on the back with 3.4×10^9 cfu *P. gingivalis* W50 approximately 3.5cm from the base of the tail.
- 5 Each day for 14 days the mice were monitored for weight loss, lesion size and behaviour. At the end of this period the mice were sacrificed as was any mouse with a weight loss greater or equal to 20% during the 14 days. At sacrifice serum was prepared from blood collected by cardiac puncture.
- 10 Lesion size data for each mouse was collected each day for 14 days and are presented in Fig 2. The lesion sizes for the two groups of animals (test and control) were analysed by ANOVA (single factor) and Mann-Whitney non-parametric statistical tests. This showed that the difference in lesion size between the test and the control groups was significant ($p=0.027$ and
- 15 $p=0.019$) using both tests examined. These results showed that the DNA vaccine construct containing the *Ag1* gene protected mice against challenge with *P. gingivalis* W50.

EXAMPLE 5.

- 20 **Immunogenicity of a synthetic peptide corresponding to the C-terminal region of Antigen 1.**

The following synthetic peptide, CIRNIWLKHKMATSAR corresponding to the C-terminal region of Ag1 was prepared based on the presence of a predicted B-cell epitope. Polyclonal antisera were raised in two dutch

25 rabbits and one New Zealand white rabbit by immunisation with the 16mer peptide conjugated to diphtheria toxoid.

- The rabbits were immunised with 78µg of peptide dissolved in TBS (50 mM Tris, 200 mM NaCl, pH 7.4) and emulsified with an equal volume of
- 30 Freund's incomplete adjuvant. The preparation was injected subcutaneously in four locations on the back. Four weeks later this was repeated and after 2 weeks a test bleed was performed and the serum response observed by ELISA against a biotinylated form of the peptide. After 2 weeks the rabbits were immunised again using the same procedures. Two weeks following the
- 35 final immunisation the rabbits were bled by cardiac puncture and the serum response determined by ELISA and western blot.

ELISA was performed using the polyclonal antisera against the biotinylated 16mer. Microtitre plates were coated with 2.54 µg/mL streptavidin. The wells were washed three times with TBS (50 mM Tris HCl pH 7.4, 200 mM NaCl) and incubated with 0.2 µg/mL biotin conjugated peptide diluted in TBS overnight at 4°C. The following day the wells were washed three times with TBS containing 0.05% v/v Tween20 (TBST) and blocked for 1 hour with 2% w/v skim milk powder at room temperature. The wells were washed three times with TBST and incubated with the polyclonal antiserum at 5-fold dilutions from 1/100 in Antibody Diluting Buffer (ADB)[TBST containing 2% w/v skim milk powder] for 2 hours at room temperature. The wells were washed three times with TBST and incubated with horseradish peroxidase conjugated goat anti-rabbit IgG diluted 1/3000 in ADB for 2 hours at room temperature. The wells were washed three times with TBST and binding of the antiserum to *P. gingivalis* proteins was visualised as above. A specific response was demonstrated indicating that specific anti-peptide antibodies had been generated. An ELISA performed using *P. gingivalis* cell-surface proteins also produced a specific response indicating that the peptide had generated antibodies capable of recognising the cell surface antigen 1 of *P. gingivalis*.

A western blot was performed to identify the components of the *P. gingivalis* cell-surface extract that was recognised by the anti-peptide antibody. After SDS-PAGE of the *P. gingivalis* cell-surface extract, the proteins were transferred to PVDF membrane and blocked as already described. The membrane was incubated with a 1/100 dilution of the antiserum in TN buffer overnight at 4°C. The following day, the membrane was washed three times in TN buffer and incubated with horseradish peroxidase conjugated goat anti-rabbit IgG diluted 1/1000 in TN buffer for 2 hours at room temperature. The membrane was washed three times in TN buffer and the binding of the serum antibodies to components of the *P. gingivalis* cell-surface protein extract was visualised as described above. A ladder of bands of 30, 33.9, 37, 40.1, 45.9, 48.8, 51.8, 56.8 and 60.3 kDa was obtained. These results indicate that the anti-peptide antibody specifically recognised Antigen 1 of *P. gingivalis* and showed that the antigen exists on the surface of *P. gingivalis* in multiple forms possibly associated with LPS.

EXAMPLE 6.

Preparation of Antigens 2, 3 and 4 (Ag2, Ag3 and Ag4).

P. gingivalis W50 was grown anaerobically at 37°C on lysed horse blood agar and in modified BM media containing 1 µg/ml hemin. Bacteria were
5 maintained on lysed horse blood plates by routine passage (< 10 passages) and used to inoculate batch cultures. Batch culture growth in Brain Heart Infusion medium was monitored at 650 nm using a spectrophotometer (295E, Perkin-Elmer). Culture purity was checked routinely by Gram stain, microscopic examination and by using a variety of biochemical tests. Stocks
10 were maintained as lyophilised cultures. A culture of *P. gingivalis* was grown to late logarithmic phase and the cells harvested by centrifugation (5,000 x g, 20 min, 4°C). Chloroform was added to the cell pellet and after gentle mixing the suspension was left for 15 min at room temperature. Following chloroform treatment, TMC buffer [20 mM Tris-HCl pH 8.0, 50
15 mM 2-mercaptoethanol and 5 mM CaCl₂] containing 50 mM NaCl was added and gently mixed. This mixture was then centrifuged (100,000 x g, 30 min, 4°C) and the supernatant filtered (0.22 µm) prior to anion-exchange FPLC. The chloroform extract was applied to an anion-exchange column (HiLoad XK 16/10 Q Sepharose, Pharmacia-LKB) cooled to 4°C, in multiple injections
20 using a 50 ml superloop (Pharmacia-LKB). The sample was eluted using a linear gradient from 0 - 100% buffer B over 90 min. The eluant was monitored at 280 nm and collected. Buffer A was TMC containing 50 mM NaCl and buffer B was TMC buffer containing 500 mM NaCl. Anion-exchange fractions were washed and then concentrated in TMC buffer
25 containing 150 mM NaCl. The fractions were then applied to a gel filtration (Superose 12) column using TMC buffer containing 150 mM NaCl. The eluant was monitored at 280 nm and peaks collected. The M_r values of eluant peaks were determined using high and low molecular mass gel filtration standards (Pharmacia-LKB). Collected peaks were then subjected
30 to reversed-phase (RP) HPLC using a C₁₈ RP-300 Brownlee column eluting with an acetonitrile gradient. The protein concentration of fractions and purified samples was determined using the Bradford protein assay (Biorad) with BSA as a standard. Chromatographic fractions and purified proteins were analysed by SDS-PAGE and Western blotting using sera from subjects
35 with *P. gingivalis* subgingivally as shown by DNA probe analysis. Serum H10 was from a subject harbouring *P. gingivalis* subgingivally but displaying no

clinical signs of periodontal disease. Serum D50 was from a patient with periodontitis who harboured *P. gingivalis* in subgingival plaque samples. From a combination of ELISA and Western blot analyses of chromatographic fractions from the anion exchange, gel filtration and reversed-phase HPLC,

- 5 three purified antigens, that were recognised by the H10 serum but were only weakly or not at all recognised by the D50 serum, were identified. The first antigen was a 30 kDa protein with an N-terminal amino acid sequence:

DNPDENPLEGDITQTHTEKYVLAED

- 10 This antigen is a fragment of a larger protein (*vide infra*) that has a molecular weight of about 46kDa, and from sequence homology is a putative fimbrial subunit protein or adhesin of *P. gingivalis*.

The second antigen was a 30 kDa protein with an N-terminal amino acid sequence:

- 15 DVLLLDVTPLSLGIETMGGVMTYLIDANTTIPKLK

This antigen, from sequence homology, is a C-terminal fragment of the *P. gingivalis* 70kDa DnaK homologue.

- 20 The third antigen recognised by the H10 serum had a molecular weight of about 10kDa and an N-terminal amino acid sequence:

VYNASISAVGNTSAIDPVVQIIHFN.

From sequence homology it is proposed that this protein is an S-layer protein of *P. gingivalis*.

25

EXAMPLE 7

Cloning and Sequence Analysis of the Gene encoding Ag3 (DnaK) of *Porphyromonas gingivalis*.

- 30 A partial BamHI *P. gingivalis* genomic library (W50 strain), constructed into bacteriophage vector LambdaGEM-12 was screened with a mixture of oligonucleotides derived from the amino acyl sequences of Ag3. The genomic library was screened by *in situ* hybridization with a mixture of degenerate oligonucleotides made to the segment of the purified protein
- 35 sequence for Ag3 the 70kDa DnaK homologue GIETMGG

Hybridisation was carried out overnight at 39°C in 6X SSC, 5X Denhardt's solution, 50µg herring sperm DNA, 1% SDS, pH 7.0. Filters were washed in 2 X SSC, 0.1% SDS at 39°C and exposed to Amersham X-ray Hyperfilm overnight at -80°C. Bacteriophage DNA was prepared from single positive
5 phage clones by the plate lysate method (See Amersham's cDNA Cloning System-λgt protocols booklet). Positively hybridizing restriction fragments were subcloned into the plasmid pUC18. Double stranded plasmid DNA was then used as sequencing template using the dideoxy-chain termination method (kits obtained from Bresatec and USB).

10

A clones containing the nucleotide sequences encoding Ag3 (DnaK clone #6) has been isolated and deposited with the Australian Government Analytical Laboratory with accession number NM/04974.

15 EXAMPLE 8

The immunogenicity of a synthetic peptide corresponding to an internal sequence of Antigen 2 (fimbrial peptide)

The peptide H-DNPDENPLEGDIT'Q'THTEKYVLAEDC-M₁ (DNP—EDC) from
20 Antigen2 (fimbrial protein) from *Porphyromonas gingivalis* strain W50 was synthesised and coupled to tetanus toxoid (TT) using 6-maleimidocaproic acyl N-hydroxysuccinimide ester forming a thioether bond. White Dutch rabbits were inoculated subcutaneously at 4 sites with 100mg of peptide coupled to the protein carrier, emulsified with incomplete Freund's adjuvant.
25 Rabbits were inoculated with a second dose on day 35 after the first injection. Rabbits were bled from the marginal ear vein on day 10 and day 84 from the first inoculation. Collected rabbit antisera was then used for ELISA and western blot analysis.

30 Detection of fimbrial peptide specific antibodies by Enzyme-Linked ImmunoSorbent Assay (ELISA).

Fimbrial peptide solution (50 µl at 5 µg ml⁻¹ of peptide) in TBS was used to coat wells of flat-bottomed polyvinyl microtiter plates (Microtiter, Dynatech Laboratories, VA., U.S.A.) overnight at room temperature in a humidified atmosphere. After removal of the coating solution, a solution (250 µl) of 1%
35 gelatin in TBS containing 0.05% Tween 20 (TBST) was added to block the remaining uncoated plastic. After 1.5 h at 37°C, the plates were washed four

times with TBST. A 1-in 100 dilution of the relevant antibodies in TBS containing 0.5% gelatin were prepared and serially diluted across the plate and incubated for 3 hours at 37°C in a humidified atmosphere. Plates were washed six times (TBST) and 50 µl of a 1/4000 dilution of horse radish peroxidase (HRPO) goat immunoglobulin (Ig) directed against rabbit Ig (BioRad, USA). After 1.5 hours at 37°C, free antibody-HRPO conjugate was removed by washing the plates six times (TBST). Bound antibody was detected by the addition of 100 µl of substrate (0.1 M acetate buffer pH 6 containing 0.01% of a 49 mM tetramethyl benzidine in DMSO and 0.004% H₂O₂). The colour reaction was stopped by the addition of 2 M H₂SO₄. Optical density (O.D.) at 455 nm was measured using a BioRad plate reader (BioRad, USA).

The binding of antisera raised against DNP—EDC-conjugate to plate bound DNP—EDC peptide was assessed. Pre-immune serum did not bind to the fimbrial peptide, however antisera from rabbits immunised with DNP—EDC-conjugate induced high titres of anti-peptide antibody. This indicates that within the peptide sequence DNP—EDC one or more B-cell epitopes are present.

20 Western blot analysis of *P. gingivalis* strain W50 cell sonicate probed with anti-DNP—EDC (fimbrial peptide) antibodies.

A SDS-Polyacrylamide gel containing low molecular weight standards (Amrad-Pharmacia, Australia), low molecular weight prestained standards (BioRad, USA) and *Porphyromonas gingivalis* W50 cell sonicate (20 µg) was prepared and transferred onto ProBlott (ABI, USA) using the following technique. ProBlott was wetted with methanol for a few seconds and placed in a tray containing the electroblotting buffer (10 mM 3-[cyclohexylamino]-1-propanesulfonic acid containing 10% methanol, pH 11). The SDS-Polyacrylamide gel of *Porphyromonas gingivalis* W50 cell sonicate was also soaked in electroblotting buffer for 5 minutes. The transblotting sandwich was assembled and electroblotted at a constant voltage of 60 volts for 90 minutes. The ProBlott was removed and rinsed with milli-Q water before staining. The ProBlott was soaked with methanol for a few seconds and the standards were stained with 0.1% Coomassie blue R-250 in 40% aqueous methanol containing 1% acetic acid. The ProBlott was destained with 50% aqueous methanol and rinsed with milli-Q water.

The ProBlott was soaked in 5% (w/v) of non-fat dry milk in 25 mM Tris/HCl in 0.5 M NaCl, pH 7.5 (TN buffer) over night at 4°C to prevent non-specific binding. The Western blot was incubated at room temperature with a 1/200 dilution of the antisera in TN buffer for 2 hours. The Western blot was then washed 4 times in TN buffer containing 0.1% triton X-100 (TNT buffer) and incubated with a 1/2000 dilution of horseradish peroxidase-conjugated goat anti-rabbit IgG (BioRad, USA) in TN buffer for 1 hour. The Western blot was then washed 4 times with TNT buffer and the protein bands detected with 0.05% 4-chloro-1-naphthol in TN buffer 16.5% ice cold methanol and 0.05% H₂O₂. Colour development was stopped with de-ionised water and the Western blot air-dried between two filter papers.

Intact fimbrial protein (adhesin) from *P. gingivalis* was detected by immunostaining using antisera raised against the fimbrial peptide-conjugate. Only two bands were detected with molecular weights of 41 and 46 kDa. This suggests that the intact fimbrial protein (adhesin) of *P. gingivalis* W50 has a molecular weight of 46 kDa and the 41 kDa band is a truncated form of the 46 kDa protein.

20 EXAMPLE 9

Preparation of Antibodies

Serum antibodies can be obtained by immunising horses, rabbits, sheep or dairy cows with Ag (1-4).

Immunizations are carried out using standard procedures. The initial immunisation is with a mixture of the antigen and Freund's incomplete adjuvant. The antibodies could be recovered from the animals serum or milk using standard procedures.

EXAMPLE 10

This example illustrates that nucleotide sequences encoding the antigens or portions thereof, can be inserted into, and expressed by various vectors including phage vectors and plasmids. Successful expression of the protein and peptides requires that either the insert comprising the gene or gene fragment, or the vector itself, contain the necessary elements for transcription and translation which is compatible with, and recognized by the particular host system used for expression. DNA encoding the antigen.

- related peptides, or oligopeptides or chimeric peptides can be synthesized or isolated and sequenced using the methods and sequences as illustrated herein. A variety of host systems may be utilized to express the antigens, related peptides or oligopeptides or chimeras, which include, but are not
- 5 limited to bacteria transformed with a bacteriophage vector, plasmid vector, or cosmid DNA; yeast containing yeast vectors; fungi containing fungal vectors; insect cell lines infected with virus (e.g. baculovirus); and mammalian cell lines transfected with plasmid or viral expression vectors, or infected with recombinant virus (e.g. vaccinia virus, adenovirus, adeno-
- 10 associated virus, retrovirus, etc.).
- Using methods known in the art of molecular biology, including methods described above, various promoters and enhancers can be incorporated into the vector or the DNA sequence encoding antigen amino acid sequences, related peptides or oligopeptide or chimeras, to increase the expression of
- 15 the antigen amino acid sequences, provided that the increased expression of the amino acid sequences is compatible with (for example, non-toxic to) the particular host cell system used. Thus and importantly, the DNA sequence can consist of the genes encoding the antigens, or any segment or combined segments of the genes which encode functional epitopes of the proteins.
- 20 Further, the DNA can be fused to DNA encoding other antigens, such as other bacterial outer membrane proteins, or other bacterial, fungal, parasitic, or viral antigens to create a genetically fused (sharing a common peptide backbone) multivalent antigen for use as an improved vaccine composition. The selection of the promoter will depend on the expression system used.
- 25 Promoters vary in strength, i.e. ability to facilitate transcription. Generally, for the purpose of expressing a cloned gene, it is desirable to use a strong promoter in order to obtain a high level of transcription of the gene and expression into gene product. For example, bacterial, phage, or plasmid promoters known in the art from which a high level of transcription have
- 30 been observed in a host cell system comprising *E. coli* include the lac promoter, trp promoter, recA promoter, ribosomal RNA promoter, the P_R and P_L promoters, lacUV5, ompF, bla, lpp, and the like, may be used to provide transcription of the inserted DNA sequence encoding antigen amino acid sequences.
- 35 Additional, if antigen, related peptides or oligopeptides or chimeras may be lethal or detrimental to the host cells, the host cell strain/line and expression

vectors may be chosen such that the action of the promoter is inhibited until specifically induced. For example, in certain operons the addition of specific inducers is necessary for efficient transcription of the inserted DNA (e.g., the lac operon is induced by the addition of lactose or isopropylthio-beta-D-galactoside). A variety of operons such as the trp operon, are under different control mechanisms. The trp operon is induced when tryptophan is absent in the growth media. The p_L promoter can be induced by an increase in temperature of host cells containing a temperature sensitive lambda repressor. In this way, greater than 95% of the promoter-directed transcription may be inhibited in uninduced cells. Thus, expression of recombinant antigenic protein, related peptides, or oligopeptides or chimeras may be controlled by culturing transformed or transfected cells under conditions such that the promoter controlling the expression from the inserted DNA encoding antigen amino acid sequences is not induced, and when the cells reach a suitable density in the growth medium, the promoter can be induced for expression from the inserted DNA.

Other control elements for efficient gene transcription or message translation include enhancers, and regulatory signals. Enhancer sequences are DNA elements that appear to increase transcriptional efficiency in a manner relatively independent of their position and orientation with respect to a nearby gene. Thus, depending on the host cell expression vector system used, an enhancer may be placed either upstream or downstream from the inserted DNA sequences encoding antigenic amino acid sequences to increase transcriptional efficiency. As illustrated previously in this example, other specific regulatory sequences have been identified which may effect the expression from the gene encoding antigen and related peptides or chimeras. These or other regulatory sites, such as transcription or translation initiation signals, can be used to regulate the expression of the gene encoding antigen, or gene fragments thereof. Such regulatory elements may be inserted into DNA sequences encoding antigenic amino acid sequences or nearby vector DNA sequences using recombinant DNA methods described herein for insertion of DNA sequences.

Accordingly, *P. gingivalis* nucleotide sequences containing regions encoding antigen, related peptides, or oligopeptides or chimeras can be ligated into an expression vector at a specific site in relation to the vector's promoter, control, and regulatory elements so that when the recombinant vector is

introduced into the host cell the *P. gingivalis*-specific DNA sequences can be expressed in the host cell. For example, the antigen-specific DNA sequences containing their own regulatory elements can be ligated into an expression vector in a relation or orientation to the vector promoter and control elements which will allow for co-expression of more than one antigen. The recombinant vector is then introduced into the appropriate host cells, and the host cells are selected, and screened for those cells containing the recombinant vector. Selection and screening may be accomplished by methods known in the art including detecting the expression of a marker gene (e.g., drug resistance marker) present in the plasmid, immunoscreening for production of specific epitopes using antisera generated to the specific antigens, and probing the DNA of the host's cells for antigen-specific nucleotide sequence using one or more oligonucleotides and methods described herein.

Genetic engineering techniques may also be used to characterize, modify and/or adapt the encoded antigenic peptides or protein. For example, site-directed mutagenesis to modify the protein in regions outside the protective domains, may be desirable to increase the safety and solubility of the subfragment to allow for easier purification and safer use. Further, genetic engineering techniques can be used to generate DNA sequences encoding a portion of the amino acid sequence of antigen. Restriction enzyme selection may be done so as not to destroy the immunopotency of the resultant peptide or oligopeptide or chimera. Antigenic sites of a protein may vary in size but can consist of from about 7 to about 14 amino acids. Thus, an antigenic protein will contain many discrete antigenic sites; therefore, many partial gene sequences could encode antigenic epitopes. Thus sequences can be constructed to contain multiple epitopes and used in an expression system to generate highly antigenic chimeric peptides or oligopeptides or proteins. Combinations of two or more proteins or peptides may result in increased immunogenicity. When using combinations of antigens these antigens may be related (*ie* from the same gene sequence or from a closely related gene from the same organism). Alternatively, the antigens may be generated from a related organism (*ie* another oral bacterium present in subgingival plaque), or from a more distantly-related organism. In particular the host organism for the vector containing the antigen-related genes and constructs can be a commensal inhabitant of the oral cavity; for example an

inhabitant of subgingival plaque, supragingival plaque or a bacterium associated with the oral mucosa. Examples of commensal intra-oral bacteria would be *Streptococcus* species and *Actinomyces* species, e.g. *Streptococcus salivarius*, *Streptococcus sanguis*, *Streptococcus gordonii*, *Actinomyces*
5 *naeslundii*. These organisms can be isolated from the periodontitis patient and then genetically engineered to express the *P. gingivalis* antigen, peptides or chimeras. The DNA encoding the antigen, peptides or chimeras could be linked with DNA encoding leader sequences of extracellular proteins of these commensal intra-oral bacteria. The DNA encoding the antigen,
10 peptides or chimeras could also be linked with, or inserted into, the DNA encoding extracellular proteins to produce secreted fusion proteins. Examples of extracellular proteins that could be used to produce fusion proteins with the antigens, peptides or chimeras could be the glucosyltransferases (GTF) or fructosyltransferases (FTF). The recombinant
15 organism would be then re-introduced into the patients oral cavity and once colonised the oral mucosa or teeth would express the *P. gingivalis* antigen, peptide, chimera or fusion to stimulate the mucosal associated lymphoid tissue to produce neutralising antibodies.

The DNA fragment encoding an antigen may be fused to other DNA
20 sequences to allow for improved expression and/or purification procedures (ie DNA sequences cloned into the vector pTrxFus, are expressed as fusions to the *E. coli* protein thioredoxin). This linkage imparts the characteristics of thioredoxin to the fusion protein which offers soluble expression of normally insoluble or difficult to express proteins. After purification, the native
25 protein is released by removal of the entire thioredoxin by digestion with enterokinase). Furthermore, the antigen may be used as a hapten by fusion to other sequences which may increase immunogenicity, if the expressed protein or peptide is not immunogenic. Alternatively, insert DNA sequences cloned into vectors may contain the native promoter sequence of the gene.
30 Therefore, transcription of the protein or peptide of interest is primarily directed from the promoter contained within the insert rather than a promoter contained within the vector sequence itself.

The plasmid expression system described uses the pUC-derived pTrcHis expression vector from Invitrogen. This vector allows high-level expression
35 of DNA sequences by the presence of the *Trc* promoter (containing the -35 region of the *Trp* promoter together with the -10 region of the *lac* promoter)

and an *rrnB* anti-terminator element. The pTrcHis vectors also contain a copy of the *lacI*^q gene which encodes the lac repressor protein. Therefore, expression of the recombinant protein/peptide is induced by addition of 1mM IPTG (de-repression) to *E. coli* grown to mid-log phase. The DNA
5 fragment is inserted into the multiple cloning site which is positioned downstream and in frame with a sequence that encodes an N-terminal fusion peptide. The N-terminal fusion peptide encodes (from 5' to 3'): an ATG translation initiation codon, a series of 6 histidine residues that function as a metal-binding domain in the translated protein, a transcript stabilising the
10 sequence from gene 10 of phage T7, and an enterokinase cleavage recognition sequence. A DNA fragment encoding a *P. gingivalis* antigen, peptide, oligopeptide or chimera is ligated to the expression vector pTrcHis. The vector was previously treated with calf intestinal phosphatase to prevent reconstitution of non-recombinant vector. The ligation mixture was used to
15 transform *E. coli* strain TOP10 by the heat-shock transformation procedure. Cells harbouring the recombinant vector plasmid were selected on Ampicillin-containing LB medium. Plasmid DNA, purified from Ampicillin-resistant colonies was analysed for the presence of recombinant insert by DNA restriction analysis and hybridisation techniques using antigen-specific
20 oligonucleotides and DNA fragments as probes. The metal-binding domain of the fusion peptide allows for one-step purification of the recombinant proteins by Immobilised Metal Affinity Chromatography. Recombinant antigen in cell culture lysates of cells harbouring the recombinant plasmid is purified by high-affinity binding to ProbondJ resin (Invitrogen). ProbondJ is
25 a nickel-charged sepharose resin that is used to purify recombinant proteins containing a poly-histidine binding domain. Bound proteins are eluted from the ProbondJ resin with either low pH buffer or by competition with imidazole or histidine. The poly-histidine leader peptide may be subsequently removed by digestion of the recombinant expressed protein
30 with Enterokinase. Enterokinase recognises the endopeptidase recognition sequence that is engineered between the poly-His affinity tag and the multiple cloning site in the vector to allow for cleavage of the poly-His tail away from the protein of interest. The purified, recombinant antigen may then be used in the generation of antibodies, vaccines and the formulation of
35 diagnostic assays as discussed.

EXAMPLE 11

Methods for using antigen-specific nucleotide sequences in molecular diagnostic assays for the detection of *P. gingivalis*. Because of the conservation of the genes encoding the antigens described herein, the nucleic acid sequences of the present invention can be used in molecular diagnostic assays for detecting *P. gingivalis* genetic material. In particular, antigen sequence-specific oligonucleotides can be synthesized for use as primers and/or probes in amplifying, and detecting amplified, nucleic acids from *P. gingivalis*. Recent advances in molecular biology have provided several means for enzymatically amplifying nucleic acid sequences. Currently the most commonly used method, PCRTM (polymerase chain reaction Cetus Corporation) involved the use of Taq Polymerase, known sequences as primers, and heating cycles which separate the replicating deoxyribonucleic acid (DNA) strands and exponentially amplify a gene of interest. Other amplification methods currently under development include LCRTM (ligase chain reaction, BioTechnica International) which utilizes DNA ligase, and a probe consisting of two halves of a DNA segment that is complementary to the sequence of the DNA to be amplified; enzyme QB replicase (Gene-Trak Systems) and a ribonucleic acid (RNA) sequence template attached to a probe complementary to the DNA to be copied which is used to make a DNA template for exponential production of complementary RNA; and NASBATM (nucleic acid sequence-based amplification, Cangene Corporation) which can be performed on RNA or DNA as the nucleic acid sequence to be amplified.

Nucleic acid probes that are capable of hybridization with specific gene sequences have been used successfully to detect specific pathogens in biological specimens at levels of sensitivity approaching 10^3 - 10^4 organisms per specimen (1990, *Gene Probes for Bacteria*, eds. Macario and deMacario, Academic Press). Coupled with a method that allows for amplification of specific target DNA sequences, species-specific nucleic acid probes can greatly increase the level of sensitivity in detecting organisms in a clinical specimen. Use of these probes may allow direct detection without relying on prior culture and/or conventional biochemical identification techniques. This embodiment of the present invention is directed to primers which amplify species-specific sequences of the genes encoding the major antigens of *P. gingivalis* described herein, and to probes which specifically hybridize

with these amplified DNA fragments. By using the nucleic acid sequences of the present invention and according to the methods of the present invention, as few as one *P. gingivalis* organism may be detected in the presence of 10 ug/ml extraneous DNA.

- 5 This embodiment is directed to species-specific oligonucleotides which can be used to amplify sequences of *P. gingivalis* DNA, if present, from DNA extracted from clinical specimens including subgingival plaque, sputum, blood, abscess and other fluids to subsequently determine if amplification has occurred. In one embodiment of the present invention, a pair of *P.*
- 10 *gingivalis*-specific DNA oligonucleotide primers are used to hybridize to *P. gingivalis* genomic DNA that may be present in DNA extracted from a clinical specimen, and to amplify the specific segment of genomic DNA between the two flanking primers using enzymatic synthesis and temperature cycling. Each pair of primers are designed to hybridize only to
- 15 the *P. gingivalis* nucleotide sequences of the present invention to which they have been synthesized to complement; one to each strand of the double-stranded DNA. Thus, the reaction is specific even in the presence of microgram quantities of heterologous DNA. For the purposes of this description, the primer derived from the sequence of the positive (gene)
- 20 strand of DNA will be referred to as the "positive primer", and the primer derived from the sequence of the negative (complementary) strand will be referred to as the "negative primer".

- Amplification of DNA may be accomplished by any one of the methods commercially available. For example, the polymerase chain reaction may be
- 25 used to amplify the DNA. Once the primers have hybridized to opposite strands of the target DNA, the temperature is raised to permit replication of the specific segment of DNA across the region between the two primers by a thermostable DNA polymerase. Then the reaction is thermocycled so that at each cycle the amount of DNA representing the sequences between the two
- 30 primers is doubled, and specific amplification of the *P. gingivalis* DNA sequences, if present, results. Further identification of the amplified DNA fragment, as being derived from *P. gingivalis* DNA, may be accomplished by liquid hybridization. This test utilizes one or more labelled oligonucleotides as probes to specifically hybridize to the amplified segment of *P. gingivalis*
- 35 DNA. Detection of the presence of sequence-specific amplified DNA may be accomplished using any one of several methods known in the art such as a

gel retardation assay with autoradiography. Thus, the nucleotide sequences of the present invention provide basis for the synthesis of oligonucleotides which have commercial applications in diagnostic kits for the detection of *P. gingivalis*. In a related embodiment, the oligonucleotides used as primers may be labeled directly, or synthesized to incorporate label. Depending on the label used, the amplification products can then be detected, after binding onto an affinity matrix, using isotopic or colorimetric detection.

DNA may be extracted from clinical specimens which may contain *P. gingivalis* using methods known in the art. For example, cells contained in the specimen may be washed in TE buffer and pelleted by centrifugation. The cells then may be resuspended in 100 ul of amplification reaction buffer containing detergents and proteinase K. Using the polymerase chain reaction, the resultant sample may be composed of the cells in 10mM Tris pH 8.3, 50mM KCl, 1.5mM MgCl₂, 0.01% gelatin, 0.45% NP40TM, 0.045% Tween 20TM, and 60 ug/ml proteinase K. The sample is incubated in a 55°C water bath for 1 hour. Following the incubation, the sample is incubated at 95°C for 10 minutes to heat-inactivate the proteinase K. The sample may then be amplified in accordance with the protocol for the polymerase chain reaction as set forth below.

The *P. gingivalis* DNA may be amplified using any one of several protocols for amplifying nucleic acids by the polymerase chain reaction. In one mode of this embodiment, the gene encoding the DnaK may be amplified from clinical isolates of *P. gingivalis* using the following conditions. DNA to be amplified (1 µg of genomic DNA) is distributed to 0.5 ml microfuge tubes and the volume adjusted to 50 ul by adding a reaction mixture comprising 0.2 mM dNTPs (dATP, dCTP, dGTP, dTTP), 0.25 ug of each positive and negative oligonucleotide primer, 1 unit of TaqI polymerase, TaqI 10x buffer (5ul), 1mM MgCl₂ (final concentration), and sterile distilled water to achieve the total volume. The TaqI polymerase is added to the reaction mixture just before use and is gently mixed, not vortexed. A layer of mineral oil, approximately 2 drops, is added to each tube and then the tubes are placed in the thermal cycler. Thirty to thirty-five cycles are general sufficient for bacterial DNA amplification. One cycle consists of 1 minute at 95°C, 1 minute at 37°C, and 1 minute at 72°C. The first cycle includes a 12 minute incubation at 95°C to assure complete denaturation.

Oligonucleotides useful as primers or probes which specifically hybridize to the gene encoding the DnaK of *P. gingivalis* and used in DNA amplification and/or detection can be biochemically synthesized, using methods known in the art, from the nucleotide sequences disclosed in the present invention.

- 5 The specificity of the oligonucleotides for *P. gingivalis* can be checked by a genebank database (Genbank) search for each individual sequence. In general, the oligonucleotides should be selected for low G-C content. For detection purposes, the oligonucleotides of the present invention may be end-labeled with a radioisotope. Probe sequences, internal to the two
- 10 primers used for amplification of the gene sequence, may be end-labeled using T4 polynucleotide kinase and gamma ^{32}P ATP. Twenty pmols of probe DNA in kinase buffer (50mM Tris, pH 7.6 10mM MgCl_2 , 5mM dithiothreitol, 0.1mM spermidine-HCl, 0.1mM EDTA, pH 8.0) is mixed with 120 uCi of gamma ^{32}P ATP and incubated at 37°C for 1 hour. Labeled probe is
- 15 separated from unincorporated label on an 8% acrylamide gel run for 1 hour at 200 volts in Tris Borate EDTA (TBE) buffer at room temperature. Labeled probe is first located by exposing the acrylamide gel to x-ray film for three minutes. The resulting autoradiogram is then positioned under the gel, and the band containing the labeled probe was excised from the gel. The gel
- 20 slice is pulverized in one milliliter of sterile distilled water, and the probe is eluted by shaker incubation overnight at 37°C. The eluted probe is separated from the gel fragments by centrifugation using a chromatography prep column. Radioactivity of the probe is determined, by counting one microliter of the labeled probe on a glass fibre filter, by liquid scintillation.
- 25 Such probe sequences may be chosen from any of the sequences disclosed herein provided the probe sequence is internal to the two primers used for amplification of the desired nucleotide sequence disclosed in the present invention.

- Alternative methods known in the art may be used to improve the detection
- 30 of amplified target sequences in accordance with the compositions and methods of the present invention. The sensitivity of detection of the amplified DNA sequences can be improved by subjecting the sequences to liquid hybridization. Alternative methods of detection known in the art, in addition to gel electrophoresis and gel electrophoresis with Southern
- 35 hybridization and autoradiography, that may be used with the compositions and methods of the present invention include: restriction enzyme digestion

with gel electrophoresis; slot-blot hybridization with a labeled oligonucleotide probe; amplification with a radiolabeled oligonucleotide probe; amplification with a radiolabeled primer with gel electrophoresis. Southern hybridization and autoradiography; amplification with a
5 radiolabeled primer with dot blot and autoradiography; amplification with oligonucleotides containing affinity tags (e.g. biotin, or one primer incorporating biotin and the other primer with a sequence specific for a DNA binding protein) followed by detection in an affinity-based assay (e.g. ELISA); and amplification with oligonucleotides containing fluorophores
10 followed by fluorescence detection.
One embodiment of non-isotopic detection involves incorporating biotin into the oligonucleotide primers of the present invention. The 5' -amino group of the primers may be biotinylated with sulfo-NHS-biotin, or biotin may be incorporated directly into the primer by synthesizing the primer in the
15 presence of biotin-labeled dNTPs. The non-isotopic labeled primers are then used in amplifying DNA from a clinical specimen. The detection for the presence or absence of amplified target sequences may be accomplished by capturing the amplified target sequences using an affinity matrix having avidin bound thereto, followed by incubation with an avidin conjugate
20 containing an enzyme which can be used to visualize the complex with subsequent substrate development. Alternatively, the amplified target sequences may be immobilized by hybridization to the corresponding probes of the target sequence wherein the probes have been affixed onto a matrix. Detection may be accomplished using an avidin conjugate containing an
25 enzyme which can be used to visualize the complex with subsequent substrate development.

EXAMPLE 12

Methods for using antigenic proteins, peptides or chimeric peptides in
30 diagnostic immunoassays.
The *P. gingivalis* antigens described herein, related peptides, oligopeptides or chimeras can be purified for use as immunogens in vaccine formulations; and as antigens for diagnostic assays or for generating *P. gingivalis*-specific antisera of therapeutic and/or diagnostic value. The antigens from *P.*
35 *gingivalis* or oligopeptides or peptides or chimeras thereof, or recombinant protein, recombinant peptides, or recombinant oligopeptides produced from

an expression vector system, can be purified with methods known in the art including detergent extraction, chromatography (e.g., ion exchange, affinity, immunoaffinity, or ultrafiltration and sizing columns), differential centrifugation, differential solubility, or other standard techniques for the purification of proteins.

In another illustration of this embodiment, a recombinant Ag1 or haeme receptor protein (HRP) was purified from a polyhistidine expression plasmid.

To purify recombinant HRP by this method, the gene encoding HRP (Fig. 1) was cloned into a polyhistidine expression vector such as plasmid pRSETA (Invitrogen Corporation), such that upon expression several histidine residues ("polyhistidine tail") are attached to the amino terminus of the HRP.

A fragment containing the gene encoding the HRP was ligated into the expression vector which had been previously restricted with an appropriate endonuclease and subsequently treated with calf intestinal phosphatase.

The ligation mixture was used to electroporate *E. coli* strain BL21(DE3) cells, and transformants were analyzed for recombinant plasmids containing the gene encoding HRP in the proper orientation with respect to the plasmid promoter. One such clone was isolated and was also shown to express the HRP when introduced into the *E. coli* host strain.

Recombinant HRP was purified as follows. A 15 ml volume of a culture of transformants containing the transformant was grown overnight in LB ampicillin broth at 37°C. The following morning, 135ml of broth was inoculated with the overnight culture and grown for 1 hour at 37°C. Cells were removed by centrifugation at 5,000 x g for 10 minutes at 4°C. The supernatant was concentrated to 10 ml by ultrafiltration through a 10kDa cut off membrane and then mixed for 10 minutes at room temperature with 1.6 ml of a resin (e.g., ProBond™, Invitrogen) which, via nickel on the resin, binds to the polyhistidine tail of the recombinant HRP. The resin was then isolated by centrifugation. The HRP was eluted from the resin by first washing the resin twice with 4 ml of denaturing wash buffer (8 M urea, 20 mM sodium phosphate, 500 mM sodium chloride, pH 7.8). The resin was then washed 2 times with 4 ml volumes of denaturing wash buffer at pH 6.0. This was followed by washing the column twice with 4 ml volumes of denaturing wash buffer at pH 4.0. Fractions of 1 ml each were collected and dialyzed against phosphate-buffered saline (PBS) containing a detergent (0.1% Triton X-100). Analysis of the eluted protein by gel electrophoresis

and Coomassie blue staining revealed a single band. It is estimated that this method results in a preparation of the protein which is 95% purified. The resultant purified recombinant HRP was immunoreactive with antibodies which recognize native protein.

- 5 As used throughout the specification, antigen oligopeptides are defined herein as a series of peptides corresponding to a portion of the amino acid sequence of the antigens described herein, as disclosed in the enclosed sequences that are synthesized as one or chemically-linked. Such peptides or oligopeptides can be synthesized using one of the several methods of
- 10 peptide synthesis known in the art including standard solid phase peptide synthesis using tertbutoxycarbonyl amino acids (Mitchell *et al.*, 1978, *J. Org. Chem.* 43:2845-2852), using 9-fluorenylmethoxycarbonyl amino acids on a polyamide support (Dryland *et al.*, 1986, *J. Chem. So. Perkin Trans. I.* 125-137); by pepscan synthesis (Geysen *et al.*, 1987, *J. Immunol. Methods* 15 03:259; 1984, *Proc. Natl. Acad. Sci. USA* 81:3998); by standard liquid phase peptide synthesis; or by recombinant expression vector systems.
- Modification of the peptides or oligopeptides, such as by deletion and substitution of amino acids (and including extensions and additions to amino acids) and in other ways, may be made so as to not substantially
- 20 detract from the immunological properties of the peptide or oligopeptide. In particular, the amino acid sequences of the antigens described herein, or peptide or oligopeptide or chimera thereof, may be altered by replacing one or more amino acids with functionally equivalent amino acids resulting in an alteration which is silent in terms of an observed difference in the
- 25 physicochemical behaviour of the protein, peptide, or oligopeptide or chimera. Functionally equivalent amino acids are known in the art as amino acids which are related and/or have similar polarity or charge. Thus, an amino acid sequence which is substantially that of the amino acid sequences depicted in the Sequence Listing herein, refers to an amino acid sequence
- 30 that contains substitutions with functionally equivalent amino acids without changing the primary biological function of protein, peptide, or oligopeptide or chimera.

In an illustration of production of chimeric peptides containing epitopes of the fimbrial protein (FP) and the HRP, defined regions of the respective

35 genes can be ligated in a construct and then expressed in an expression system wherein the plasmid expression vector (pGEX2T) directs the

synthesis of foreign polypeptides in *E. coli* as a fusion peptides with glutathione-S-transferase (GST), a 26 kilodalton protein from *Schistosoma japonicum*. In this mode of the embodiment, and using *P. gingivalis* DNA as the template, selected regions of the genes encoding FP-HRP can be amplified by the polymerase chain reaction using selected oligonucleotides. The oligonucleotides are designed so that the resulting amplified gene fragments after ligation contains a BamHI restriction site on its 5' end, and an EcoRI restriction site on the 3' end so that the amplified chimeric fragment can be directionally cloned into pGEX2T. The sequence of each recombinant plasmid is confirmed by dideoxy sequencing. To purify each recombinant peptide, the respective recombinant plasmid containing the chimeric gene fragment is transformed into *E. coli* JM109. The transformant is grown in 400ml of LB broth containing 25ug/ml of ampicillin by adding forty ml of an overnight culture to 360 ml of broth, and incubating for 1.5 hours at 37°C with shaking. IPTG is added to 0.01mM and the culture incubated for an additional 3 hours. Cells are centrifuged at 5000 x g and the cell pellet resuspended in 5 ml of PBS. Cells are sonicated and the mixture centrifuged at 10,000 x g for 10 minutes. The supernatant is mixed with 0.5 ml of preswelled glutathione-agarose beads. After mixing for 2 minutes at room temperature, the beads (with fusion peptide bound to the glutathione) are washed 2 additional times with PBS containing 1% triton-X-100. The beads are then washed once in 0.05M Tris, pH 8.0. To cleave the FP-HRP chimeric peptide from the glutathione-S-transferase, the washed beads are incubated in 0.25% (final concentration) human thrombin in Tris buffer for 1 hour at room temperature. A protease inhibitor, PMSF, is then added to a concentration of 100 ug/ml. The beads are removed by centrifugation and the supernatant contains the purified FP-HRP chimeric peptide. Immunoblot assays affirm that the fusion peptides are immunoreactive, with Ag1- and Ag2-specific rabbit polyclonal antisera

Purified antigens, peptides, oligopeptides and chimeras may be used as antigens in immunoassays for the detection of *P. gingivalis*-specific antisera present in the body fluid of an individual suspected of having an infection caused by *P. gingivalis*. The detection of Ag(1-4) or related peptides in immunoassays, includes any immunoassay known in the art including, but not limited to, radioimmunoassay, enzyme-linked immunosorbent assay

(ELISA), "sandwich" assay, precipitin reaction, agglutination assay, fluorescent immunoassay, and chemiluminescence-based immunoassay.

EXAMPLE 13

- 5 Methods and compounds for formulations related to Ag(1-4) and related peptides and chimeras.

This embodiment of the present invention relates to antigens and/or peptides or oligopeptides or chimeras thereof, to be used in as immunogens in a prophylactic and/or therapeutic compositions for active immunization to
10 protect against or treat infections caused by *P. gingivalis*. For vaccine purposes, an antigen of *P. gingivalis* comprising a bacterial protein should be immunogenic, and induce functional antibodies directed to one or more surface-exposed epitopes on intact bacteria, wherein the epitope(s) are conserved amongst strains of *P. gingivalis*.

- 15 In one illustration of the invention, Ag4 having the properties desirable of a vaccine antigen, the protein can be purified from *P. gingivalis* using the method described herein in Example 6. Mice are immunized with the purified Ag4 protein (25 ug) with adjuvant (20 ug of QS21) two times at four week intervals. Blood from the immunized mice is drawn 32 days after the
20 last immunization and the immune sera was pooled. The pooled immune sera assayed against whole bacteria (*P. gingivalis* strain W50) by an enzyme linked immunosorbent assay (ELISA). For the whole cell ELISA, overnight cultures of bacteria are harvested by a swab and suspended in PBS to an absorbance of 0.1 at 600nm. Aliquots (100 ul) of the bacterial suspension are
25 added to the wells of a 96 well microtiter plate and dried overnight at room temperature. The plates are blocked with 100ul of 0.1% (w/v) gelatin in PBS. This, and all remaining incubations, are for one hour at room temperature. The blocking solution is removed and 100 ul of the immune sera, diluted in PBS with 0.1% (w/v) gelatin, is added to the wells and incubated. After
30 washing three times with PBS, the bound antibodies are detected by incubating with 100 ul of alkaline phosphatase conjugated recombinant protein G (1:1500 in PBS with 0.1% (w/v) gelatin). The plates are washed and color development is facilitated by the addition of 100 ul/well of p-nitrophenyl phosphate (2 mg/ml in diethanolamine). after 30 minutes, the
35 reaction is stopped by adding 50 ul of 3M NaOH. The absorbance is read at 492 nm using an ELISA reader. Endpoint titers are determined as the

reciprocal of the dilution at which the absorbance is greater than that of the blank wells

In further illustrating that antigens from W50 possesses properties desirable of a vaccine antigen, pooled immune sera raised to strain W50 was shown to have cross-reactivity with heterologous strains.

For vaccine development, antigen-specific amino acid sequences may be purified from a host containing a recombinant vector which expresses antigen or related peptides or chimeras. Such hosts include, but are not limited to, bacterial transformants, yeast transformants, filamentous fungal transformants, and cultured cells that have been either infected or transfected with a vector which encodes antigen amino acid sequences. Peptides or oligopeptides or chimeras corresponding to portions of the antigens may be produced from chemical or enzymatic cleavage of the antigens; or chemically synthesized using methods known in the art and with the amino acid sequence deduced from the nucleotide sequence of the genes encoding the antigens as a reference. Alternatively, peptides may be produced from a recombinant vector. The protein, peptide, or oligopeptide or chimera immunogen is included as the relevant immunogenic material in the vaccine formulation, and in therapeutically effective amounts, to induce an immune response. Many methods are known for the introduction of a vaccine formulation into the human or animal to be vaccinated. These include, but are not limited to, intradermal, intramuscular, intraperitoneal, intravenous, subcutaneous, ocular, intranasal, and oral administration. The vaccine may further comprise a physiological carrier such as a solution, a polymer or liposomes; and an adjuvant, or a combination thereof. Various adjuvants are used in conjunction with vaccine formulations. The adjuvants aid by modulating the immune response and in attaining a more durable and higher level of immunity using smaller amounts of vaccine antigen or fewer doses than if the vaccine antigen were administered alone. Examples of adjuvants include incomplete Freund's adjuvant (IFA), Adjuvant 65 (containing peanut oil, mannide monooleate and aluminium monostearate), oil emulsions, Ribi adjuvant, the pluronic polyols, polyamines, Avridine, Quil A, saponin, MPL, QS-21, and mineral gels such as aluminium salts. Other examples include oil in water emulsions such as SAF-1, SAF-0, MF59, Seppic ISA720, and other particulate adjuvants such as ISCOMs and ISCOM matrix. An extensive but not exhaustive list of other examples of adjuvants

are listed in Cox and Coulter 1992 [In : Wong WK (ed.) *Animals parasite control utilising technology*. Boca Raton: CRC press. 1992: 49-112] In addition to the adjuvant the vaccine may include conventional pharmaceutically acceptable carriers, excipients, fillers, buffers or diluents as appropriate. One or more doses of the vaccine containing adjuvant may be administered prophylactically to prevent periodontitis or therapeutically to treat already present periodontitis.

Another embodiment of this mode of the invention involves the production of antigen-specific amino acid sequences as a hapten, i.e. a molecule which cannot by itself elicit an immune response. In such case, the hapten may be covalently bound to a carrier or other immunogenic molecule which will confer immunogenicity to the coupled hapten when exposed to the immune system. Thus, such a antigen-specific hapten linked to a carrier molecule may be the immunogen in a vaccine formulation.

Another mode of this embodiment provides for either a live recombinant viral vaccine, recombinant bacterial vaccine, recombinant attenuated bacterial vaccine, or an inactivated recombinant viral vaccine which is used to protect against infections caused by *P. gingivalis*. Vaccinia virus is the best known example, in the art, of an infectious virus that is engineered to express vaccine antigens derived from other organisms. The recombinant live vaccinia virus, which is attenuated or otherwise treated so that it does not cause disease by itself, is used to immunize the host. Subsequent replication of the recombinant virus within the host provides a continual stimulation of the immune system with the vaccine antigens such as Ag(1-4) protein, related peptides or chimeras, thereby providing long lasting immunity.

Other live vaccine vectors include: adenovirus, cytomegalovirus, and preferably the poxviruses such as vaccinia (Paoletti and Panicali, U.S. Patent No. 4,603,112) and attenuated *Salmonella* strains (Stocker *et al.*, U.S. Patent Nos. 5,210,035; 4,837,151; and 4,735,801; and Curtiss *et al.*, 1988, Vaccine 6:155-160). Live vaccines are particularly advantageous because they continually stimulate the immune system which can confer substantially long-lasting immunity. When the immune response is protective against subsequent *P. gingivalis* infection, the live vaccine itself may be used in a preventive vaccine against *P. gingivalis*. In particular, the live vaccine can be based on a bacterium that is a commensal inhabitant of the oral cavity

This bacterium can be transformed with a vector carrying a recombinant Ag(1-4), peptides, oligopeptides or chimeric peptides and then used to colonise the oral cavity, in particular the oral mucosa. Once colonised the oral mucosa, the expression of the recombinant protein, peptide or chimera will stimulate the mucosal associated lymphoid tissue to produce neutralising antibodies.

To further illustrate this mode of the embodiment, using molecular biological techniques such as those illustrated in Example 12, the genes encoding Ag(1-4) or gene fragments encoding one or more peptides or chimeras may be inserted into the vaccinia virus genomic DNA at a site which allows for expression of epitopes but does not negatively affect the growth or replication of the vaccinia virus vector. The resultant recombinant virus can be used as the immunogen in a vaccine formulation. The same methods can be used to construct an inactivated recombinant viral vaccine formulation except that the recombinant virus is inactivated, such as by chemical means known in the art, prior to use as an immunogen and without substantially affecting the immunogenicity of the expressed immunogen. A mixture of inactivated viruses which express different epitopes may be used in the formulation of a multivalent inactivated vaccine. In either case, the inactivated recombinant vaccine or mixture of inactivated viruses may be formulated with a suitable adjuvant in order to enhance the immunological response to the vaccine antigens.

In another variation of this embodiment, genetic material is used directly as the vaccine formulation. Nucleic acid (DNA or RNA) containing sequences encoding Ag(1-4), related peptides or oligopeptides or chimeras, operatively linked to one or more regulatory elements can be introduced directly to vaccinate the individual ("direct gene transfer") against pathogenic strains of *P. gingivalis*. Direct gene transfer into a vaccinated individual, resulting in expression of the genetic material by the vaccinated individual's cells such as vascular endothelial cells as well as the tissue of the major organs, has been demonstrated by techniques in the art such as by injecting intravenously an expression plasmid:cationic liposome complex (Zhu *et al.*, 1993, *Science* 261:209-211). Other effective methods for delivering vector DNA into a target cell are known in the art. In one example, purified recombinant plasmid DNA containing viral genes has been used to inoculate (whether parentally, mucosally, or via gene-gun immunization) vaccines to

- induce a protective immune response (Fynan *et al.*, 1993, *Proc. Natl. Acad. Sci. USA* 90:11478-11482). In another example, cells removed from an individual can be transfected or electroporated by standard procedures known in the art, resulting in the introduction of the recombinant vector DNA into the target cell. Cells containing the recombinant vector DNA may then be selected for using methods known in the art such as via a selection marker expressed in the vector, and the selected cells may then be re-introduced into the individual to express the antigenic protein, related peptides or oligopeptides or chimeras.
- One preferred method of vaccination with genetic material comprises the step of administering to the individual the nucleic acid molecule that comprises a nucleic acid sequence that encodes for one or more of Ag(1-4), related peptides, or oligopeptides or chimeras, wherein the nucleic acid molecule is operatively linked to one or more regulatory sequences necessary for expression. The nucleic acid molecule can be administered directly, or first introduced into a viral vector and administered via the vector. The nucleic acid molecule can be administered in a pharmaceutically acceptable carrier or diluent and may contain compounds that can enhance the effectiveness of the vaccine. These additional compounds include, but are not limited to, adjuvants that enhance the immune response, and compounds that are directed to modulate the immune response, e.g. cytokines, collectively referred to as "immune modulators"; or other compounds which increase the uptake of nucleic acid by the cells, referred to as "nucleic acid uptake enhancers". The immunization with the nucleic acid molecule can be through any parental route (intravenous, intraperitoneal, intradermal, subcutaneous, or intramuscular), or via contact with mucosal surfaces of the nasopharynx, trachea, or gastrointestinal tract.
- As an alternative to active immunization, immunization may be passive, i.e. immunization comprising administration of purified immunoglobulin containing antibody against Ag(1-4) epitopes.

EXAMPLE 14

- The following is an example of a proposed toothpaste formulation containing anti-Ag1 antibodies

Ingredient

w/w

49

Dicalcium phosphate dihydrate	50.0
Glycerol	20.0
Sodium carboxymethyl cellulose	1.0
Sodium lauryl sulphate	1.5
Sodium lauroyl sarconisate	0.5
Flavour	1.0
Sodium saccharin	0.1
Chlorhexidine gluconate	0.01
Dextranase	0.01
Goat serum containing anti-Ag1	0.2
Water	balance

EXAMPLE 15

The following is an example of a proposed toothpaste formulation

<u>Ingredient</u>	<u>% w/w</u>
Dicalcium phosphate dihydrate	50.0
Sorbitol	10.0
Glycerol	10.0
Sodium carboxymethyl cellulose	1.0
Sodium lauryl sulphate	1.5
Sodium lauroyl sarcosinate	0.5
Flavour	1.0
Sodium saccharin	0.1
Sodium monofluorophosphate	0.3
Chlorhexidine gluconate	0.01
Dextranase	0.01
Bovine serum containing anti-Ag2	0.2
Water	balance

EXAMPLE 16

5 The following is an example of a proposed toothpaste formulation.

<u>Ingredient</u>	<u>% w/w</u>
Dicalcium phosphate dihydrate	50.0
Sorbitol	10.0
Glycerol	10.0
Sodium carboxymethyl cellulose	1.0
Lauroyl diethanolamide	1.0
Sucrose monolaurate	2.0
Flavour	1.0
Sodium saccharin	0.1
Sodium monofluorophosphate	0.3
Chlorhexidine gluconate	0.01
Dextranase	0.01
Bovine milk Ig containing anti-Ag1	0.1
Water	balance

EXAMPLE 17

The following is an example of a proposed toothpaste formulation.

<u>Ingredient</u>	<u>% w/w</u>
Sorbitol	22.0
Irish moss	1.0
Sodium Hydroxide (50%)	1.0
Gantrez	19.0
Water (deionised)	2.69
Sodium Monofluorophosphate	0.76
Sodium saccharine	0.3
Pyrophosphate	2.0
Hydrated alumina	48.0
Flavour oil	0.95
anti-Ag1 mouse monoclonal	0.3
sodium lauryl sulphate	2.00

EXAMPLE 18

- 5 The following is an example of a proposed liquid toothpaste formulation.

<u>Ingredient</u>	<u>% w/w</u>
Sodium polyacrylate	50.0
Sorbitol	10.0
Glycerol	20.0
Flavour	1.0
Sodium saccharin	0.1
Sodium monofluorophosphate	0.3
Chlorhexidine gluconate	0.01
Ethanol	3.0
Equine Ig containing anti-Ag2	0.2
Linolic acid	0.05
Water	balance

EXAMPLE 19

The following is an example of a proposed mouthwash formulation.

<u>Ingredient</u>	<u>% w/w</u>
Ethanol	20.0
Flavour	1.0
Sodium saccharin	0.1
Sodium monofluorophosphate	0.3
Chlorhexidine gluconate	0.01
Lauroyl diethanolamide	0.3
Rabbit Ig containing anti-Ag1	0.2
Water	balance

EXAMPLE 20

- 5 The following is an example of a proposed mouthwash formulation.

<u>Ingredient</u>	<u>% w/w</u>
Gantrez S-97.	2.5
Glycerine	10.0
Flavour oil	0.4
Sodium monofluorophosphate	0.05
Chlorhexidine gluconate	0.01
Lauroyl diethanolamide	0.2
Mouse anti-Ag2 monoclonal	0.3
Water	balance

EXAMPLE 21

The following is an example of a proposed lozenge formulation.

<u>Ingredient</u>	<u>% w/w</u>
Sugar	75-80
Corn syrup	1-20
Flavour oil	1-2
NaF	0.01-0.05
Mouse anti-Ag1 monoclonal	0.3
Mg stearate	1-5
Water	balance

EXAMPLE 22

- 5 The following is an example of a proposed gingival massage cream formulation.

<u>Ingredient</u>	<u>% w/w</u>
White petrolatum	8.0
Propylene glycol	4.0
Stearyl alcohol	8.0
Polyethylene Glycol 4000	25.0
Polyethylene Glycol 400	37.0
Sucrose monostearate	0.5
Chlorohexidine gluconate	0.1
Mouse anti-Ag2 monoclonal	0.3
Water	balance

EXAMPLE 23

The following is an example of a proposed chewing gum formulation

<u>Ingredient</u>	<u>% w/w</u>
Gum base	30.0
Calcium carbonate	2.0
Crystalline sorbitol	53.0
Glycerine	0.5
Flavour oil	0.1
Mouse anti-Ag1 and anti-Ag2 monoclonals	0.3
Water	balance

- 5 It will be appreciated by persons skilled in the art that numerous variations and/or modifications may be made to the invention as shown in the specific embodiments without departing from the spirit or scope of the invention as broadly described. The present embodiments are, therefore, to be considered in all respects as illustrative and not restrictive.

CLAIMS

- 1 A composition for use in raising an immune response directed against *Porphyromonas gingivalis*, the composition including a suitable adjuvant and/or acceptable carrier and one substantially purified *P. gingivalis* immunogen, the immunogen being selected from the group consisting of Antigen 1, Antigen 2, Antigen 3, Antigen 4 and epitope containing fragments thereof, in which:-

Antigen 1 is an antigen of *P. gingivalis* and has an internal amino acid sequence:

10 DLENKGEATLLVTFGSSYKAPRETYAKIEKTFAAAYPDQR;

Antigen 2 is an antigen of *P. gingivalis* and has an internal amino acid sequence:

DNPDENPLEGDITQTHTEKYVLAED

15 Antigen 3 is an antigen of *P. gingivalis* and has an internal amino acid sequence:

DVLLLDVTPLSLGIETMGGVMTYLIDANTTIPKLLK

Antigen 4 is an antigen of *P. gingivalis* and has an internal amino acid sequence:

VYNASISAVGNTSAIDPVVQIIHHN.

- 20 2. A composition as claimed in claim 1 in which the composition further includes at least one additional purified *P. gingivalis* immunogen, the immunogen being selected from the group consisting of Antigen 1, Antigen 2, Antigen 3, Antigen 4 and epitope containing fragments thereof, in which:-

25 Antigen 1 is an antigen of *P. gingivalis* and has an internal amino acid sequence:

DLENKGEATLLVTFGSSYKAPRETYAKIEKTFAAAYPDQR;

Antigen 2 is an antigen of *P. gingivalis* and has an internal amino acid sequence:

DNPDENPLEGDITQTHTEKYVLAED

30 Antigen 3 is an antigen of *P. gingivalis* and has an internal amino acid sequence:

DVLLLDVTPLSLGIETMGGVMTYLIDANTTIPKLLK

Antigen 4 is an antigen of *P. gingivalis* and has an N-terminal amino acid sequence:

35 VYNASISAVGNTSAIDPVVQIIHHN.

- 3 A composition as claimed in claim 1 or claim 2 in which Antigen 1 has an amino acid sequence as shown in Fig 1.
- 4 A composition as claimed in claim 1 or claim 2 in which Antigen 3 includes an amino acid sequence encoded by the open reading frame of the
- 5 clone deposited with AGAL under accession No. NM 97/04974 which hybridises with degenerate probes corresponding to the amino acid sequence DVLLLDVTPLSLGIETMGGVMTYLIDANTTIPKLLK.
5. A substantially purified *P. gingivalis* antigen or epitope containing fragment thereof, wherein antigen has an internal amino acid sequence:
- 10 DLENKGEATLLVTFGSSYKAPRETYAKIEKTFAAAYPDQR.
6. A substantially purified *P. gingivalis* antigen or epitope containing fragment thereof as claimed in claim 5 in which the antigen has an amino acid sequence as shown in Fig 1.
7. A substantially purified *P. gingivalis* antigen or epitope containing
- 15 fragment thereof, wherein antigen has an internal amino acid sequence. DNPDENPLEGDITQTHTEKYVLAED.
8. A substantially purified *P. gingivalis* antigen or epitope containing fragment thereof, wherein antigen has an internal amino acid sequence:
- 20 DVLLLDVTPLSLGIETMGGVMTYLIDANTTIPKLLK.
9. A substantially purified *P. gingivalis* antigen or epitope containing fragment thereof as claimed in claim 8 in which the antigen includes an amino acid sequence encoded by the open reading frame of the clone deposited with AGAL under accession No. NM 97/04974 which hybridises with degenerate probes corresponding to the amino acid sequence
- 25 DVLLLDVTPLSLGIETMGGVMTYLIDANTTIPKLLK.
10. A substantially purified *P. gingivalis* antigen or epitope containing fragment thereof, wherein antigen has an internal amino acid sequence:
- VYNASISAVGNTSAIDPVVQIIHHN.
- 30 11 An antibody preparation comprising antibodies raised against the composition as claimed in any one of claims 1 to 4 or against the antigen as claimed in any one of claims 5 to 10.
- 12 An antibody preparation as claimed in claim 11 in which the antibodies are polyclonal antibodies.
- 35 13 An antibody preparation as claimed in claim 11 in which the antibodies are monoclonal antibodies.

- 14 A method of treating a subject suffering from *Porphyromonas gingivalis* infection, the method comprising administering to the subject an amount of the antibody preparation as claimed in any one of claims 11 to 13 effective to at least partially suppress the pathogenic effects of
- 5 *Porphyromonas gingivalis*.
- 15 A method as claimed in claim 14 in which the antibody preparation is administered as a mouth wash or as a dentifrice.
- 16 A method of reducing the prospect of *P. gingivalis* infection in an individual and/or severity of disease, the method comprising administering
- 10 to the individual an amount of the composition as claimed in any one of claims 1 to 4 effective to induce an immune response in the individual directed against *P. gingivalis*.
- 17 A composition for use in raising an immune response directed against *Porphyromonas gingivalis*, the composition including a suitable
- 15 adjuvant and/or acceptable carrier and a DNA molecule including a sequence encoding one *P. gingivalis* immunogen, the immunogen being selected from the group consisting of Antigen 1, Antigen 2, Antigen 3, Antigen 4 and epitope containing fragments thereof, in which:-
- Antigen 1 is an antigen of *P. gingivalis* and has an internal amino acid sequence:
- 20 DLENKGEATLLVTFGSSYKAPRETYAKIEKTFAAAYPDQR;
- Antigen 2 is an antigen of *P. gingivalis* and has an internal amino acid sequence:
- DNPDENPLEGDITQTHTEKYVLAED
- 25 Antigen 3 is an antigen of *P. gingivalis* and has an internal amino acid sequence:
- DVLLLDVTPLSLGIETMGGVMITYLIDANTTIPKLLK
- Antigen 4 is an antigen of *P. gingivalis* and has an internal amino acid sequence:
- 30 VYNASISAVGNTSAIDPVVQIIHHN.
- 18 A composition as claimed in claim 17 in which the DNA molecule includes a sequence as shown in Fig 1 or an epitope encoding fragment thereof.
- 19 A composition as claimed in claim 17 in which the DNA molecule
- 35 includes a sequence as present in the open reading frame of the clone deposited with AGAL under accession No. NM 97/04974 which hybridises

with degenerate probes corresponding to the amino acid sequence
DVLLLDVTPLSLGIETMGGVMTYLIDANTTIPKLIK, or an epitope encoding
fragment thereof

20. An isolated DNA molecule, the DNA molecule having a sequence
5 encoding the amino acid sequence shown in Fig 1, or an epitope thereof.
21. An isolated DNA molecule as claimed in claim 20 in which the DNA
molecule has the sequence shown in Fig 1 or an epitope encoding fragment
thereof.
22. An isolated DNA molecule, the DNA molecule having a sequence as
10 present in the open reading frame of the clone deposited with AGAL under
accession No. NM 97/04974 which hybridises with degenerate probes
corresponding to the amino acid sequence
DVLLLDVTPLSLGIETMGGVMTYLIDANTTIPKLIK.
23. A nucleotide probe, the probe having a sequence which hybridises
15 specifically to the DNA molecule as claimed in any one of claims 20 to 22.
24. A polynucleotide probe as claimed in claim 23 in which the probe is
labelled.
25. A recombinant host cell, the host cell being transformed with a DNA
sequence(s) encoding at least one *P. gingivalis* immunogen operatively
20 linked to control sequences such that under appropriate conditions the host
cell expresses the at least one immunogen, the immunogen being selected
from the group consisting of Antigen 1, Antigen 2, Antigen 3, Antigen 4 and
epitope containing fragments thereof, in which:-
Antigen 1 is an antigen of *P. gingivalis* and has an internal amino
25 acid sequence:
DLENKGEATLLVTFGSSYKAPRETYAKIEKTFAAAYPDQR;
Antigen 2 is an antigen of *P. gingivalis* and has an internal amino
acid sequence:
DNPDENPLEGDITQTHTEKYVLAED
30 Antigen 3 is an antigen of *P. gingivalis* and has an internal amino
acid sequence:
DVLLLDVTPLSLGIETMGGVMTYLIDANTTIPKLIK
Antigen 4 is an antigen of *P. gingivalis* and has an internal amino
acid sequence:
35 VYNASISAVGNTSAIDPVVQIIHHN.

26 A recombinant host cell as claimed in claim 25 in which the host cell is an oral commensal.

1 TTGGGACTCGCCACATCGCATCGTTTCGGCTCATTTCGGCCACATCATCTCGATCGAAGG
61 CTGGGTATAGACAACCGTTTGACTACGTGGATGATCGTCCTATGGGTGTCAATTATGGTA
121 CCGTAACGGCGGGACGTACTTTCTTTGCTCAAATAGCGATTGCGATTCAACAACCTAATGTC
181 TCACAAATTAATTTAAGAACAGAGATGAAAAAAGTGAATTTAGCGACTTTGGGACTTATG
M K K L I L A T L G L M
241 GCCATTGCCATGCTCTCATGTTCAAGCAACAACAAGGATTGAGAGAACAAGGGGAGGCT
A I A M L S C S S N N K D L E N K G E A
301 ACTCTTTTGGTAACGTTTGGTAGCTCCTATAAAGCTCCACGCGAAACCTATGCCAAGATT
T L L V T F G S S Y K A P R E T Y A K I
361 GAGAAGACTTTTGGCCGACGTTTATCCCGATCAAAGGATAAGCTGGACATACAGCTCTTCT
E K T F A A A Y P D Q R I S W T Y T S S
421 ATTATCCGAAAGAACTGGCTCAGCAGGGTATTTATATCGATGCTCCGGATGAGGCTTTC
I I R K K L A Q Q G I Y I D A P D E A L
481 GAGAAATTGGCTCGTCTGGGTTATAAGAAGATCAATGTACAGAGTCTTCATGTGATTCCC
E K L A R L G Y K K I N V Q S L H V I P
541 GGCCGAGAATATGATGAGATGATCGACTTTGTCATAAGTTTAAAGGCAGCACATAGTCAAT
G R E Y D E M I D F V N K F K A A H S E
601 ATTACTGTGAAGGTAGGGCGTCCGCTTTTCGATACCGATGAAGATATGGCGAGGTGGCA
I T V K V G R P L F D T D E D M R E V A
661 GAGATCTTGCACAGCGTTTTCAGCAACGATAGAGAAAGGTGAAGCTATTGTATTTCATG
E I L H K R F Q Q T I E K G E A I V F M
721 GGACACGGCACCAGCATGCTGCCAATGACAGGTATGCCCGTATCAATAAGATCATGAAG
G H G T E H A A N D R Y A R I N K I M K
781 AACTATAGCAAGTTCATGATCGTCGGAACCGTCGAGTCCGATCGCTTATCAATGATGTT
N Y S K F M I V G T V E S D P S I N D V
841 ATTGCCGAACCTGAAAGAAACCGGTGCCACGGCCGTAACAATGATGCCGCTGATGAGTGT
I A E L K E T G A T A V T M M P L M S V
901 GCAGGCGACCATGCTACGAATGATATGGCCGGAGATGAGGACGATAGCTGGAAGACGTTG
A G D H A T N D M A G D E D D S W K T L
961 CTGACCAATGCCGGCTACACAGTTTCTATAGACAAGCTGGACAATGGCAATTTCTCAGCT
L T N A G Y T V S I D K L D N G N F S A
1021 CTTGGAGATATAGAAGAGATCCGGAATATCTGGCTCAAGCATATGAAAGCCACCTCTGCT
L G D I E E I R N I W L K H M K A T S A
1081 CGCTAAGGACGGGCGSATATGCAATGAGACAATCAAGCAATTAGGTTACGAGAGCACTTA
R

FIGURE 1

2 / 2




FIGURE 2

INTERNATIONAL SEARCH REPORT

International Application No.
PCT/AU 97/00212

A. CLASSIFICATION OF SUBJECT MATTER																						
Int Cl ⁶ C07K 14:195, 15:12; C12N 001/21; 15/31; G01N 033/569; A61K 39/02; 39/40																						
According to International Patent Classification (IPC) or to both national classification and IPC																						
B. FIELDS SEARCHED																						
Minimum documentation searched (classification system followed by classification symbols)																						
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched JAPIO																						
Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) WPAT (Porphyromonas, and Bacteroides and gingivalis) SYN: ANGIS; USPM																						
C. DOCUMENTS CONSIDERED TO BE RELEVANT																						
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.																				
E	US 5523390 (Travis, J. et al); June 4, 1996, Claims 1 and 2; page 1.	1-26																				
E	WO 96/17936 (University of Florida); June 13, 1996, Claims 1-20	1-26																				
A	AU-B- 21148/92 (Eastman Kodak Company)	1-26																				
<input checked="" type="checkbox"/> Further documents are listed in the continuation of Box C <input checked="" type="checkbox"/> See patent family annex																						
<p>* Special categories of cited documents:</p> <table border="0"> <tr> <td>"A"</td> <td>document defining the general state of the art which is not considered to be of particular relevance</td> <td>"T"</td> <td>later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</td> </tr> <tr> <td>"E"</td> <td>earlier document but published on or after the international filing date</td> <td>"X"</td> <td>document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone</td> </tr> <tr> <td>"L"</td> <td>document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</td> <td>"Y"</td> <td>document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art</td> </tr> <tr> <td>"O"</td> <td>document referring to an oral disclosure, use, exhibition or other means</td> <td>"&"</td> <td>document member of the same patent family</td> </tr> <tr> <td>"P"</td> <td>document published prior to the international filing date but later than the priority date claimed</td> <td></td> <td></td> </tr> </table>			"A"	document defining the general state of the art which is not considered to be of particular relevance	"T"	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention	"E"	earlier document but published on or after the international filing date	"X"	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone	"L"	document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"Y"	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art	"O"	document referring to an oral disclosure, use, exhibition or other means	"&"	document member of the same patent family	"P"	document published prior to the international filing date but later than the priority date claimed		
"A"	document defining the general state of the art which is not considered to be of particular relevance	"T"	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention																			
"E"	earlier document but published on or after the international filing date	"X"	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone																			
"L"	document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"Y"	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art																			
"O"	document referring to an oral disclosure, use, exhibition or other means	"&"	document member of the same patent family																			
"P"	document published prior to the international filing date but later than the priority date claimed																					
Date of the actual completion of the international search 21 April 1997		Date of mailing of the international search report 29 APR 1997																				
Name and mailing address of the ISA/AU AUSTRALIAN INDUSTRIAL PROPERTY ORGANISATION PO BOX 200 WOODEN ACT 2606 AUSTRALIA Facsimile No. (06) 283 3929		Authorized officer Madhu K. Joga Telephone No. (06) 283 2512																				

INTERNATIONAL SEARCH REPORT

International Application No.
PCT/AU 97/00212

C (Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	April 22, 1993. Claims 1-11	1-16
	WO 95/26404 (Kyowa Hakko Koggo et al) October 5, 1995	17-26

INTERNATIONAL SEARCH REPORT
Information on patent family members

International Application No.
PCT/AU 97/00212

This Annex lists the known "A" publication level patent family members relating to the patent documents cited in the above-mentioned international search report. The Australian Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

Patent Document Cited in Search Report			Patent Family Member			
US	5523390	AU	21148/92	WO	9617936	WO 9526404
	WO 9507286		EP 537828		AU45151/96	
	EP 717747		US 5399484			
						END OF ANNEX